

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
**WO 02/088123 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 471/04**,  
A61K 31/435, A61P 15/10

(74) Agent: **NAPOLI, James, J.**; Marshall, Gerstein & Borun,  
6300 Sears Tower, 233 South Wacker Drive, Chicago, IL  
60606 (US).

(21) International Application Number: PCT/US02/10367

(22) International Filing Date: 2 April 2002 (02.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/286,730 25 April 2001 (25.04.2001) US

(71) Applicant (for all designated States except US): **LILLY  
ICOS LLC** [US/US]; 1209 Orange Street, Wilmington, DE  
19801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SAWYER, Jason,  
Scott** [US/US]; 5718 North Winthrop Avenue, Indianapo-  
lis, In 46220 (US). **ORME, Mark, W.** [US/US]; 4235  
Francis Avenue #203, Seattle, WA 98103 (US). **COPP,  
James, D.** [US/US]; 1430 Runyon Road, Greenwood, IN  
46143 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GI,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

Published:


— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 02/088123 A1

(54) Title: CHEMICAL COMPOUNDS



(I)

(57) Abstract: Compounds of general structural formula (I) and use of the compounds and salts and solvates thereof, as therapeutic agents as inhibitors of phosphodiesterase 5 cardiovascular disorders.

- 1 -


## CHEMICAL COMPOUNDS

FIELD AND BACKGROUND OF THE INVENTION

5 This invention relates to a series of compounds, to methods of preparing the compounds, to pharmaceutical compositions containing the compounds, and to their use as therapeutic agents. In particular, the invention relates to compounds that  
 10 are potent and selective inhibitors of cyclic guanosine 3',5'-monophosphate specific phosphodiesterase (cGMP-specific PDE), in particular PDE5, and have utility in a variety of therapeutic areas wherein such inhibition is considered beneficial, including  
 15 the treatment of cardiovascular disorders and erectile dysfunction.

SUMMARY OF THE INVENTION

20 The present invention provides compounds of formula (I)




30 wherein  $R^0$ , independently, is selected from the group consisting of halo,  $C_{1-6}$ alkyl, aryl, heteroaryl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{3-8}$ cycloalkylQ,  $C(=O)R^a$ ,  $OC(=O)R^a$ ,  $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ ,

- 2 -

$C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkyleneC(=O)OR<sup>a</sup>, C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>c</sup>,  
 C(=O) $C_{1-4}$ alkyleneHet, C(=O)NR<sup>a</sup>R<sup>b</sup>, C(=O)NR<sup>a</sup>R<sup>c</sup>, C(=O)-  
 NR<sup>a</sup> $C_{1-4}$ alkyleneOR<sup>b</sup>, C(=O)NR<sup>a</sup> $C_{1-4}$ alkyleneHet, OR<sup>a</sup>, OC<sub>1-4</sub>-  
 alkyleneC(=O)OR<sup>a</sup>, OC<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, OC<sub>1-4</sub>alkyleneHet,  
 5 OC<sub>1-4</sub>alkyleneOR<sup>a</sup>, OC<sub>1-4</sub>alkyleneNR<sup>a</sup>C(=O)OR<sup>b</sup>, NR<sup>a</sup>R<sup>b</sup>,  
 NR<sup>a</sup> $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>, NR<sup>a</sup>C(=O)R<sup>b</sup>, NR<sup>a</sup>C(=O)NR<sup>a</sup>R<sup>b</sup>,  
 N(SO<sub>2</sub> $C_{1-4}$ alkyl)<sub>2</sub>, NR<sup>a</sup>(SO<sub>2</sub> $C_{1-4}$ alkyl), nitro, trifluoro-  
 methyl, trifluoromethoxy, cyano, SO<sub>2</sub>NR<sup>a</sup>R<sup>b</sup>, SO<sub>2</sub>R<sup>a</sup>, SOR<sup>a</sup>,  
 SR<sup>a</sup>, and OSO<sub>2</sub>CF<sub>3</sub>;

10 R<sup>1</sup> is selected from the group consisting of  
 optionally substituted aryl, optionally substituted  
 heteroaryl, an optionally substituted C<sub>3-8</sub>cycloalkyl  
 ring, an optionally substituted C<sub>3-8</sub>heterocycloalkyl  
 ring, an optionally substituted bicyclic ring


15




20

wherein the fused ring A is a 5- or 6-membered ring,  
 saturated or partially or fully unsaturated, and  
 comprises carbon atoms and optionally one to three  
 25 heteroatoms selected from oxygen, sulfur, and  
 nitrogen, hydrogen, C<sub>1-6</sub>alkyl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>-  
 alkylenearyl, haloC<sub>1-6</sub>alkyl, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,  
 C<sub>1-4</sub>alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkenyl,  
 C<sub>3-8</sub>heterocycloalkenyl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkylene-  
 30 QR<sup>a</sup>, C<sub>2-6</sub>alkenyleneQR<sup>a</sup>, C<sub>1-4</sub>alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>,


- 3 -



5




10



15

and a spiro substituent having a structure



25

;

30  $R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{2-6}$ alkenyl,  $C_{1-3}$ alkylenearyl, aryl $C_{1-3}$ alkyl,  $C(=O)R^a$ , aryl, heteroaryl,  $C(=O)R^a$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $SO_2R^a$ ,  $SO_2NR^aR^b$ ,  $S(=O)R^a$ ,

- 4 -

S(=O)NR<sup>a</sup>R<sup>b</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyl-  
eneHet, C(=O)C<sub>1-4</sub>alkylenearyl, C(=O)C<sub>1-4</sub>alkylene-  
heteroaryl, C<sub>1-4</sub>alkylenearyl substituted with one or  
more of SO<sub>2</sub>NR<sup>a</sup>R<sup>b</sup>, NR<sup>a</sup>R<sup>b</sup>, C(=O)OR<sup>a</sup>, NR<sup>a</sup>SO<sub>2</sub>CF<sub>3</sub>, CN, NO<sub>2</sub>,  
5 C(=O)R<sup>a</sup>, OR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, and OC<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>,  
C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkyleneC-  
(=O)C<sub>1-4</sub>alkylenearyl, C<sub>1-4</sub>alkyleneC(=O)C<sub>1-4</sub>alkylene-  
heteroaryl, C<sub>1-4</sub>alkyleneC(=O)Het, C<sub>1-4</sub>alkyleneC(=O)-  
NR<sup>a</sup>R<sup>b</sup>, C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>C(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyl-  
10 eneOC<sub>1-4</sub>alkyleneOR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, C<sub>1-4</sub>alkyleneC-  
(=O)OR<sup>a</sup>, and C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>;

R<sup>3</sup> is selected from the group consisting of  
C(=O)R<sup>b</sup>, C(=O)OR<sup>b</sup>, C(=O)NR<sup>a</sup>R<sup>b</sup>, C(=O)NR<sup>a</sup>R<sup>c</sup>, C(=S)NR<sup>a</sup>R<sup>b</sup>,  
C(=S)NR<sup>a</sup>R<sup>c</sup>, C(=O)Het, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOR<sup>a</sup>,  
15 C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet, C(=O)C<sub>1-4</sub>alkylenearyl,  
C(=O)C<sub>1-4</sub>alkyleneheteroaryl, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkylenearyl,  
C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneC<sub>3-8</sub>cycloalkyl, C(=O)NR<sup>b</sup>SO<sub>2</sub>R<sup>c</sup>,  
C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOC<sub>1-6</sub>alkyl, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkylene-  
heteroaryl, NR<sup>a</sup>R<sup>c</sup>, NR<sup>a</sup>C(=O)R<sup>b</sup>, NR<sup>a</sup>C(=O)NR<sup>a</sup>R<sup>c</sup>,  
20 NR<sup>a</sup>(SO<sub>2</sub>C<sub>1-4</sub>alkyl), N(SO<sub>2</sub>C<sub>1-4</sub>alkyl)<sub>2</sub>, OR<sup>a</sup>,  
NR<sup>a</sup>C(=O)C<sub>1-4</sub>alkyleneN(R<sup>b</sup>)<sub>2</sub>, NR<sup>a</sup>C(=O)C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,  
NR<sup>a</sup>(C(=O)C<sub>1-3</sub>alkylenearyl, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkylene-  
C<sub>3-8</sub>heterocycloalkyl, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkyleneHet, and  
C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>b</sup>;


25 R<sup>4</sup> is selected from the group consisting of  
hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl,  
C<sub>1-3</sub>alkylenearyl, C<sub>1-3</sub>alkyleneHet, C<sub>3-8</sub>cycloalkyl, and  
C<sub>3-8</sub>heterocycloalkyl;

X is selected from the group consisting of  
30 C(=O), (CH<sub>2</sub>)<sub>t</sub>C(=O), C(=O)C≡C, C(=O)C(R<sup>a</sup>)=C(R<sup>a</sup>), C(=S),  
SO, SO<sub>2</sub>, SO<sub>2</sub>C(R<sup>a</sup>)=CR<sup>a</sup>, CR<sup>a</sup>R<sup>b</sup>, CR<sup>a</sup>=CR<sup>a</sup>, C(=O)NR<sup>a</sup>, and  
C(=N-OR<sup>a</sup>);

- 5 -

Y is selected from the group consisting of  
 $R^a$ ,  $R^d$ ,  $(CH_2)_n C(=O)R^c$ ,  $N(R^b)(CH_2)_n R^c$ ,  $O(CH_2)_n R^c$ ,  
 $N(R^b)C(=O)R^c$ ,  $C(=O)N(R^a)(R^c)$ ,  $N(R^a)C(=O)R^c$ ,

5



10

and  $N(R^a)SO_2R^c$ ;

$R^a$  is selected from the group consisting of  
hydrogen,  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl, aryl,  
aryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenearyl, heteroaryl,  
15 heteroaryl $C_{1-3}$ alkyl, and  $C_{1-3}$ alkyleneheteroaryl;

$R^b$  is selected from the group consisting of  
hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{1-3}$ alkylene $N(R^a)_2$ ,  
aryl, aryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenearyl, and heteroaryl;

$R^c$  is selected from the group consisting of  
20 hydrogen,  $C_{1-6}$ alkyl, aryl, heteroaryl, aryl $C_{1-3}$ alkyl,  
heteroaryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylene $N(R^a)_2$ ,  $C_{1-6}$ alkylene-  
aryl,  $C_{1-6}$ alkyleneHet, halo $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  
 $C_{3-8}$ heterocycloalkyl, Het,  $C_{1-3}$ alkyleneheteroaryl,  
 $C_{1-6}$ alkylene $C(=O)OR^a$ , and  $C_{1-3}$ alkylene $C_{3-8}$ heterocyclo-  
25 alkyl;

or  $R^a$  and  $R^c$  are taken together to form a  
5- or 6-membered ring, optionally containing at  
least one heteroatom;

$R^d$  is a 5- or 6-membered ring or a bicyclic  
30 fused ring system, saturated or partially or fully  
unsaturated, comprising carbon atoms and optionally  
one to three heteroatoms selected from oxygen,

- 6 -

sulfur, and nitrogen, and optionally substituted  
with one or more R<sup>e</sup> or R<sup>f</sup>;

R<sup>e</sup> is selected from the group consisting of

nitro,

5

trifluoromethyl,

trifluoromethoxy,

halogen,

cyano,

Het,

10

C<sub>1-6</sub>alkyl,

C<sub>1-6</sub>alkyleneOR<sup>a</sup>,

C(=O)R<sup>a</sup>,

OC(=O)R<sup>a</sup>,

C(=O)OR<sup>a</sup>,

15

C<sub>1-4</sub>alkyleneHet,

C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,

OC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,

C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,

C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>f</sup>,

20

C(=O)C<sub>1-4</sub>alkyleneHet,

C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>,

C<sub>2-6</sub>alkenyleneNR<sup>a</sup>R<sup>b</sup>,

C(=O)NR<sup>a</sup>R<sup>b</sup>,

C(=O)NR<sup>a</sup>R<sup>g</sup>,

25

C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOR<sup>b</sup>,

C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet,

OR<sup>a</sup>,

OC<sub>2-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>,

OC<sub>1-4</sub>alkyleneCH(OR<sup>a</sup>)CH<sub>2</sub>-NR<sup>a</sup>R<sup>b</sup>,

30

OC<sub>1-4</sub>alkyleneHet,

OC<sub>2-4</sub>alkyleneOR<sup>a</sup>,

OC<sub>2-4</sub>alkyleneNR<sup>a</sup>C(=O)OR<sup>b</sup>,

NR<sup>a</sup>R<sup>b</sup>,

- 7 -

 $\text{NR}^a\text{C}_{1-4}\text{alkyleneNR}^a\text{R}^b$ , $\text{NR}^a\text{C}(=\text{O})\text{R}^b$ , $\text{NR}^a\text{C}(=\text{O})\text{NR}^a\text{R}^b$ , $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,5  $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ , $\text{SO}_2\text{NR}^a\text{R}^b$ ,and  $\text{OSO}_2\text{trifluoromethyl}$ ;

$\text{R}^f$  is selected from the group consisting of hydrogen, halogen,  $\text{OR}^a$ ,  $\text{C}_{1-6}\text{alkyl}$ , nitro, and  $\text{NR}^a\text{R}^b$ ;

10 or  $\text{R}^e$  and  $\text{R}^f$  are taken together to form a 3- or 4-membered alkylene or alkenylene chain component of a 5- or 6-membered ring, optionally containing at least one heteroatom;

15  $\text{R}^g$  is phenyl or  $\text{C}_{4-6}\text{cycloalkyl}$ , optionally substituted with one or more halogen,  $\text{C}(=\text{O})\text{OR}^a$ , or  $\text{OR}^a$ ;

 $\text{Q}$  is O, S, or  $\text{NR}^h$ ; $\text{B}$  is O, S, or  $\text{NR}^h$ ; $\text{C}$  is O, S, or  $\text{NR}^a$ ;20  $\text{D}$  is  $\text{CR}^a$  or N; $\text{E}$  is  $\text{CR}^a$ ,  $\text{C}(\text{R}^a)_2$ , or  $\text{NR}^h$ ; and

25  $\text{R}^h$  is null or is selected from the group consisting of hydrogen,  $\text{C}_{1-6}\text{alkyl}$ , aryl, heteroaryl,  $\text{arylC}_{1-3}\text{alkyl}$ , heteroaryl $\text{C}_{1-3}\text{alkyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ , and  $\text{C}_{1-3}\text{alkyleneheteroaryl}$ ;

30 Het represents a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with  $\text{C}_{1-4}\text{alkyl}$  or  $\text{C}(=\text{O})\text{OR}^a$ ;

 $n$  is 0 or 1; $q$  is 0, 1, 2, 3, or 4;




- 8 -

t is 1, 2, 3, or 4;

and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

The present invention also provides compounds of structural formula (Ia), i.e., compounds of structural formula (I) wherein  $(X)_n$ -Y is hydrogen, i.e., a compound having a formula



15


wherein  $R^0$ , independently, is selected from the group consisting of halo,  $C_{1-6}$ alkyl, aryl, heteroaryl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{3-8}$ cycloalkylQ,  $C(=O)R^a$ ,  $OC(=O)R^a$ ,  $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^c$ ,  $C(=O)C_{1-4}$ alkyleneHet,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $NR^aR^b$ ,  $OC_{1-4}$ alkyleneHet,  $OC_{1-4}$ alkylene $OR^a$ ,  $OC_{1-4}$ alkylene $NR^aC(=O)OR^b$ ,  $NR^aR^b$ ,  $NR^aC_{1-4}$ alkylene $NR^aR^b$ ,  $NR^aC(=O)R^b$ ,  $NR^aC(=O)NR^aR^b$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $NR^a(SO_2C_{1-4}alkyl)$ , nitro, trifluoromethyl, trifluoromethoxy, cyano,  $SO_2NR^aR^b$ ,  $SO_2R^a$ ,  $SOR^a$ ,  $SR^a$ , and  $OSO_2CF_3$ ;

$R^1$  is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, an optionally substituted  $C_{3-8}$ cycloalkyl

- 9 -


ring, an optionally substituted C<sub>3-8</sub>heterocycloalkyl  
 ring, an optionally substituted bicyclic ring

5




10 wherein the fused ring A is a 5- or 6-membered ring,  
 saturated or partially or fully unsaturated, and  
 comprises carbon atoms and optionally one to three  
 heteroatoms selected from oxygen, sulfur, and  
 nitrogen, hydrogen, C<sub>1-6</sub>alkyl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>-  
 15 alkylenearyl, haloC<sub>1-6</sub>alkyl, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,  
 C<sub>1-4</sub>alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkenyl,  
 C<sub>3-8</sub>heterocycloalkenyl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkylene-  
 QR<sup>a</sup>, C<sub>2-6</sub>alkenyleneQR<sup>a</sup>, C<sub>1-4</sub>alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>,

20




25




30

- 10 -



5

and a spiro substituent having a structure



10

15

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{2-6}$ alkenyl,  $C_{1-3}$ alkylenearyl, aryl $C_{1-3}$ alkyl,  $C(=O)R^a$ , aryl, heteroaryl,  $C(=O)R^a$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $SO_2R^a$ ,  $SO_2NR^aR^b$ ,  $S(=O)R^a$ ,  $S(=O)NR^aR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneOR<sup>a</sup>,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)C_{1-4}$ alkylenearyl,  $C(=O)C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkylenearyl substituted with one or more of  $SO_2NR^aR^b$ ,  $NR^aR^b$ ,  $C(=O)OR^a$ ,  $NR^aSO_2CF_3$ , CN,  $NO_2$ ,  $C(=O)R^a$ ,  $OR^a$ ,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>, and  $OC_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneC(=O)Het,  $C_{1-4}$ alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>C(=O)R<sup>a</sup>,  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneC(=O)OR<sup>a</sup>, and  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>;

30

- 11 -

$R^3$  is selected from the group consisting of  
 $C(=O)R^b$ ,  $C(=O)OR^b$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  
 $C(=S)NR^aR^c$ ,  $C(=O)Het$ ,  $C(=O)NR^aC_{1-4}alkyleneOR^a$ ,  $C(=O)-$   
 $NR^aC_{1-4}alkyleneHet$ ,  $C(=O)C_{1-4}alkylenearyl$ ,  $C(=O)C_{1-4}-$   
 $alkyleneheteroaryl$ ,  $C(=O)NR^aC_{1-4}alkylenearyl$ ,  $C(=O)-$   
 $NR^aC_{1-4}alkyleneC_{3-8}cycloalkyl$ ,  $C(=O)NR^bSO_2R^c$ ,  $C(=O)-$   
 $NR^aC_{1-4}alkyleneOC_{1-6}alkyl$ ,  $C(=O)NR^aC_{1-4}alkylenehetero-$   
 $aryl$ ,  $NR^aR^c$ ,  $NR^aC(=O)R^b$ ,  $NR^aC(=O)NR^aR^c$ ,  $NR^a(SO_2C_{1-4}-$   
 $alkyl)$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $OR^a$ ,  $NR^aC(=O)C_{1-4}alkylene-$   
 $N(R^b)_2$ ,  $NR^aC(=O)C_{1-4}alkyleneC(=O)OR^a$ ,  $NR^aC(=O)C_{1-3}-$   
 $alkylenearyl$ ,  $NR^aC(=O)C_{1-3}alkyleneC_{3-8}heterocycloalkyl$ ,  
 $NR^aC(=O)C_{1-3}alkyleneHet$ , and  $C(=O)NR^aSO_2R^b$ ;

$R^4$  is selected from the group consisting of  
hydrogen,  $C_{1-6}alkyl$ ,  $aryl$ ,  $heteroaryl$ ,  $arylC_{1-3}alkyl$ ,  
 $C_{1-3}alkylenearyl$ ,  $C_{1-3}alkyleneHet$ ,  $C_{3-8}cycloalkyl$ , and  
 $C_{3-8}heterocycloalkyl$ ;

$R^a$  is selected from the group consisting of  
hydrogen,  $C_{1-6}alkyl$ ,  $C_{3-6}cycloalkyl$ ,  $aryl$ ,  
 $arylC_{1-3}alkyl$ ,  $C_{1-3}alkylenearyl$ ,  $heteroaryl$ ,  
 $heteroarylC_{1-3}alkyl$ , and  $C_{1-3}alkyleneheteroaryl$ ;

$R^b$  is selected from the group consisting of  
hydrogen,  $C_{1-6}alkyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{1-3}alkyleneN(R^a)_2$ ,  
 $aryl$ ,  $arylC_{1-3}alkyl$ ,  $C_{1-3}alkylenearyl$ , and  $heteroaryl$ ;

$R^c$  is selected from the group consisting of  
hydrogen,  $C_{1-6}alkyl$ ,  $aryl$ ,  $heteroaryl$ ,  $arylC_{1-3}alkyl$ ,  
 $heteroarylC_{1-3}alkyl$ ,  $C_{1-3}alkyleneN(R^a)_2$ ,  $C_{1-6}alkylene-$   
 $aryl$ ,  $C_{1-6}alkyleneHet$ ,  $haloC_{1-6}alkyl$ ,  $C_{3-8}cycloalkyl$ ,  
 $C_{3-8}heterocycloalkyl$ ,  $Het$ ,  $C_{1-3}alkyleneheteroaryl$ ,  
 $C_{1-6}alkyleneC(=O)OR^a$ , and  $C_{1-3}alkyleneC_{3-8}heterocyclo-$   
 $alkyl$ ;

or  $R^a$  and  $R^c$  are taken together to form a  
5- or 6-membered ring, optionally containing at  
least one heteroatom;

- 12 -

Q is O, S, or  $\text{NR}^h$ ;

B is O, S, or  $\text{NR}^h$ ;

C is O, S, or  $\text{NR}^a$ ;

D is  $\text{CR}^a$  or N;

5 E is  $\text{CR}^a$ ,  $\text{C}(\text{R}^a)_2$ , or  $\text{NR}^h$ ; and

$\text{R}^h$  is null or is selected from the group consisting of hydrogen,  $\text{C}_{1-6}$ alkyl, aryl, heteroaryl, aryl $\text{C}_{1-3}$ alkyl, heteroaryl $\text{C}_{1-3}$ alkyl,  $\text{C}_{1-3}$ alkylenearyl, and  $\text{C}_{1-3}$ alkyleneheteroaryl;

10 Het represents a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted  
15 with  $\text{C}_{1-4}$ alkyl or  $\text{C}(=\text{O})\text{OR}^a$ ;

n is 0 or 1;

q is 0, 1, 2, 3, or 4;

t is 1, 2, 3, or 4;

20 and pharmaceutically acceptable salts and solvates thereof.

As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, propyl, and butyl groups.  
25 The hydrocarbon group can contain up to 16 carbon atoms. The term "alkyl" also includes "bridged alkyl," i.e., a  $\text{C}_4$ - $\text{C}_{16}$  bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo-  
30 [3.2.1]octyl, or decahydronaphthyl. The term "cycloalkyl" is defined as a cyclic  $\text{C}_3$ - $\text{C}_8$  hydrocarbon group, e.g., cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl.

- 13 -

5 The term "alkenyl" is defined identically as "alkyl," except for containing a carbon-carbon double bond. "Cycloalkenyl" is defined similarly to cycloalkyl, except a carbon-carbon double bond is present in the ring.

10 The term "alkylene" refers to an alkyl group having a substituent. For example, the term "C<sub>1-3</sub>alkylenearyl" refers to an alkyl group containing one to three carbon atoms, and substituted with an aryl group. The term "alkenylene" as used herein is similarly defined, and contains the indicated number of carbon atoms and a carbon-carbon double bond, and includes straight chained and branched alkenylene groups, like ethylenylene.

15 The term "halo" or "halogen" is defined herein to include fluorine, bromine, chlorine, and iodine.

20 The term "haloalkyl" is defined herein as an alkyl group substituted with one or more halo substituents, either fluoro, chloro, bromo, iodo, or combinations thereof. Similarly, "halocycloalkyl" is defined as a cycloalkyl group having one or more halo substituents.

25 The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl. Unless otherwise indicated, an "aryl" group can be unsubstituted or substituted, for example, with one or  
30 more, and in particular one to three, halo, alkyl, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Exemplary aryl groups

- 14 -

include phenyl, naphthyl, tetrahydronaphthyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 4-methoxyphenyl, 3-trifluoromethylphenyl, 4-nitrophenyl, and the like. The terms  
5 "arylC<sub>1-3</sub>alkyl" and "heteroarylC<sub>1-3</sub>alkyl" are defined as an aryl or heteroaryl group having a C<sub>1-3</sub>alkyl substituent.

The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one  
10 or two aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring, and which can be unsubstituted or substituted, for example, with one or more, and in particular one to three, substituents, like halo, alkyl, hydroxy,  
15 hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Examples of heteroaryl groups include thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazolyl, isothiazolyl, isoxazolyl, imidizolyl, benzothiazolyl,  
20 pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "heterocycloalkyl" is defined as monocyclic, bicyclic, and tricyclic groups containing one or more, e.g., one to three, heteroatoms  
25 independently selected from the group consisting of oxygen, nitrogen, and sulfur. A "heterocycloalkyl" group also can contain an oxo group (=O) attached to the ring. Nonlimiting examples of heterocycloalkyl groups include 1,3-dioxolane, 2-pyrazoline,  
30 pyrazolidine, pyrrolidine, piperazine, a pyrroline, 2H-pyran, 4H-pyran, morpholine, thiopholine, piperidine, 1,4-dithiane, and 1,4-dioxane.

The term "hydroxy" is defined as -OH.

- 15 -

The term "alkoxy" is defined as -OR, wherein R is alkyl.

The term "alkoxyalkyl" is defined as an alkyl group wherein a hydrogen has been replaced by an alkoxy group. The term "(alkylthio)alkyl" is defined similarly as alkoxyalkyl, except a sulfur atom, rather than an oxygen atom, is present.

The term "hydroxyalkyl" is defined as a hydroxy group appended to an alkyl group.

The term "amino" is defined as -NH<sub>2</sub>, and the term "alkylamino" is defined as -NR<sub>2</sub>, wherein at least one R is alkyl and the second R is alkyl or hydrogen.

The term "acylamino" is defined as RC(=O)N, wherein R is alkyl or aryl.

The term "alkylthio" is defined as -SR, wherein R is alkyl.

The term "alkylsulfinyl" is defined as R-SO<sub>2</sub>, wherein R is alkyl.

The term "alkylsulfonyl" is defined as R-SO<sub>3</sub>, wherein R is alkyl.

The term "nitro" is defined as -NO<sub>2</sub>.

The term "trifluoromethyl" is defined as -CF<sub>3</sub>.

The term "trifluoromethoxy" is defined as -OCF<sub>3</sub>.

The term "spiro" as used herein refers to a group having two carbon atoms directly bonded to the carbon atom to which R<sup>1</sup> is attached.

The term "cyano" is defined as -CN.

In a preferred embodiment, q is 0. In other preferred embodiments, R<sup>0</sup> is selected from the group consisting of aryl, Het, OR<sup>a</sup>, C(=O)OR<sup>a</sup>, C<sub>1-4</sub>-




- 16 -

alkyleneNR<sup>a</sup>R<sup>b</sup>, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkylQ, C(=O)NR<sup>a</sup>R<sup>b</sup>, and C(=O)NR<sup>a</sup>R<sup>c</sup>.

In a preferred group of compounds of formula (I), R<sup>1</sup> is represented by


5



10

wherein the bicyclic ring can represent, for example, naphthalene or indene, or a heterocycle, such as benzoxazole, benzothiazole, benzisoxazole, benzimidazole, quinoline, indole, benzothiophene, or benzofuran, or

15



20


wherein p is an integer 1 or 2, and G, independently, is C(R<sup>a</sup>)<sub>2</sub>, O, S, or NR<sup>a</sup>. The bicyclic ring comprising the R<sup>1</sup> substituent typically is attached to the rest of the molecule by a phenyl ring carbon atom.

25

In another preferred group of compounds of formula (I), R<sup>1</sup> is represented by an optionally substituted bicyclic ring

30

- 17 -




5


wherein  $p$  is 1 or 2, and  $G$ , independently, are  $C(R^a)_2$  or  $O$ .

Especially preferred  $R^1$  substituents  
include


10



15




20



25


- 18 -



5


,

10




15

,




20

,



25

, and



30





.

Within this particular group of compounds, nonlimiting examples of substituents for the bicyclic ring

- 19 -

include halogen (e.g., chlorine), C<sub>1-3</sub>alkyl (e.g., methyl, ethyl, or i-propyl), OR<sup>a</sup> (e.g., methoxy, ethoxy, or hydroxy), CO<sub>2</sub>R<sup>a</sup>, halomethyl or halomethoxy (e.g., trifluoromethyl or trifluoromethoxy), cyano, nitro, and NR<sup>a</sup>R<sup>b</sup>.


In other preferred embodiments, R<sup>1</sup> is optionally substituted and selected from the group consisting of C<sub>1-4</sub>alkyleneQR<sup>a</sup>, C<sub>1-4</sub>alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkenyl, C<sub>1-6</sub>alkyl,




- 20 -

In a more preferred group of compounds of formula (I),  $R^1$  is represented by


5



10



15




$C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkenyl,  $C_{1-6}$ alkyl,  $C_{1-4}$ alkyleneQR<sup>a</sup>, and  $C_{1-4}$ alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>. A preferred Q is oxygen.


20

Some preferred  $R^1$  substituents are


25




30




- 21 -



5




10



15


 $-\text{CH}_2\text{OR}^a$ ,  $-\text{CH}_2\text{OCH}_2\text{OR}^a$ ,

20



, and

25



30


- 22 -

Within this particular group of compounds, preferred  $R^a$  substituents include hydrogen,  $C_{1-6}$ alkyl, and benzyl.

In a preferred embodiment,  $R^2$  is selected from the group consisting of hydrogen, aryl, hetero-  
 5 aryl,  $OR^a$ ,  $NR^aR^b$ ,  $NR^aR^c$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene-  
 heteroaryl,  $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ -  
 alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $OR^a$ ,  $C_{1-4}$ alkyleneC(=O)-  
 $NR^aR^b$ ,  $C_{1-4}$ alkyleneC(=O) $NR^aR^c$ ,  $C_{1-4}$ alkyleneC(=O)Het,  
 10  $C_{1-4}$ alkylene $NR^aR^b$ ,  $C_{1-4}$ alkylene $NR^aR^c$ ,  $C_{1-4}$ alkylene $NR^aC$ -  
 (=O) $R^a$ , and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ .


In more preferred embodiments,  $R^2$  is selected from the group consisting of hydrogen;  $C_{1-4}$ -  
 alkyleneheteroaryl, wherein the heteroaryl group is  
 15 selected from the group consisting of benzimidazole,  
 a triazole, and imidazole;  $C_{1-4}$ alkyleneHet, wherein  
 Het is selected from the group consisting of piper-  
 azine, morpholine, pyrrolidine, pyrrolidone, tetra-  
 hydrofuran, piperidine,

20



, and

25




30

;

- 23 -

$C_{1-4}$ alkylene $C_6H_5$ , optionally substituted with one to three groups selected from the group consisting of  $C(=O)OR^a$ ,  $NR^aR^b$ ,  $NR^aSO_2CF_3$ ,  $SO_2NR^aR^b$ ,  $CN$ ,  $OR^a$ ,  $C(=O)R^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ , nitro,  $OC_{1-4}$ alkylenearyl, and  $OC_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)$ benzyl;  $C_{1-4}$ alkylene $C(=O)OR^a$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^c$ ;  $C_{1-4}$ alkyleneHet;  $NR^aR^b$ ;  $OH$ ;  $OC_{1-4}$ alkyl;  $C_6H_5$ ;  $C_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $OR^a$ ;  $C_{1-4}$ alkylene $NHC(=O)R^a$ ; and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ . In most preferred embodiments,  $R^2$  is hydrogen.

In preferred embodiments,  $R^3$  is selected from the group consisting of  $C(=O)OR^a$ ,  $C(=O)R^a$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OC_{1-6}$ alkyl,  $C(=O)NR^aC_{1-4}$ alkylene- $C_{3-8}$ cycloalkyl,  $C(=O)Het$ ,



$C(=O)NR^aC_{1-4}$ alkyleneheteroaryl,  $C(=O)NR^aC_{1-4}$ alkylenearyl,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)NR^aR^c$ , and  $C(=S)NR^aR^c$ ;

In preferred embodiments,  $R^4$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, and heteroaryl.

In preferred embodiments,  $n$  is 0, or  $X$  is selected from the group consisting of  $C(=O)$ ,  $C(=O)CR^a=CR^a$ ,  $(CH_2)_tC=O$ ,  $C(=S)$ , and  $C(=N-OR^a)$ .


In preferred embodiments,  $Y$  is selected from the group consisting of  $R^a$ ,  $R^d$ ,  $N(R^b)(CH_2)_nR^c$ ,  $O(CH_2)_nR^c$ ,  $N(R^b)C(=O)R^c$ , and  $N(R^a)C(=O)R^c$ .




- 24 -

In especially preferred embodiments, q is  
0 or R<sup>0</sup> is selected from the group consisting of  
halo, methyl, trifluoromethyl, and trifluoromethyl;  
R<sup>1</sup> is selected from the group consisting of


5



10




15



20


25




30

- 25 -


5



10




15




20

25



, and

- 26 -




5

;


$R^2$  is selected from the group consisting of hydrogen,  
 $C_{1-6}$ alkyl,  $C(=O)NR^aR^c$ , and  $C_{1-4}$ alkyleneHet;  $R^3$  is  
 10 selected from the group consisting of  $C(=O)OC_2H_5$ ,  
 $C(=O)OCH_3$ ,  $C(=O)NHCH_2C_6H_5$ ,  $C(=O)NH(CH_2)_2C_6H_5$ ,  
 $C(=O)NHC_6H_5$ ,  $C(=O)NH_2$ ,  $C(=O)N(CH_3)_2$ ,  $C(=S)N(CH_3)_2$ ,  
 $C(=O)NH(CH_2)_2CH_3$ ,  $C(=O)N(CH_2)_3CH_3$ ,  $C(=O)NHCH_3$ ,  
 $C(=O)NHCH(CH_3)_2$ ,  $C(=O)NH(CH_2)_3OCH_3$ ,

15




20

,



25


,



30


,

- 27 -




5

,




10

,



15


,



20

25


,




30

,


- 28 -




5




10




15



20




25




30


- 29 -




5



10




15



20


25



30

, and

- 30 -




5

;

$R^4$  is selected from the group consisting of hydrogen and  $C_{1-6}$ alkyl; X is selected from the group consisting of  $C(=O)$ ,  $C(=S)$ ,  $CH_2C(=O)$ , and  $C(=O)CH=CH$ , or n is 0; and Y is selected from the group consisting of  $NH(CH_2)_3CH_3$ ,  $OCH_3$ ,  $C_6H_5$ , and


15



20

An especially preferred subclass of compounds within the general scope of formula (I) is represented by compounds of formula (II)

25



30

(II)

- 31 -

and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

Compounds of formula (I) can contain one or more asymmetric center, and, therefore, can exist as stereoisomers. The present invention includes both mixtures and separate individual stereoisomers of the compounds of formula (I). Compounds of formula (I) also can exist in tautomeric forms, and the invention includes both mixtures and separate individual tautomers thereof.

Pharmaceutically acceptable salts of the compounds of formula (I) can be acid addition salts formed with pharmaceutically acceptable acids. Examples of suitable salts include, but are not limited to, the hydrochloride, hydrobromide, sulfate, bisulfate, phosphate, hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzenesulfonate, and p-toluenesulfonate salts. The compounds of formula (I) also can provide pharmaceutically acceptable metal salts, in particular alkali metal salts and alkaline earth metal salts, with bases. Examples include the sodium, potassium, magnesium, and calcium salts.

Compounds of the present invention are potent and selective inhibitors of cGMP-specific PDE5. Thus, compounds of formula (I) are of interest for use in therapy, specifically for the treatment of a variety of conditions where selective inhibition of PDE5 is considered to be beneficial.

Phosphodiesterases (PDEs) catalyze the hydrolysis of cyclic nucleotides, such as cyclic adenosine monophosphate (cAMP) and cyclic guanosine



- 32 -

monophosphate (cGMP). The PDEs have been classified into at least seven isoenzyme families and are present in many tissues (J.A. Beavo, *Physiol. Rev.*, 75, p. 725 (1995)).

5 PDE5 inhibition is a particularly attractive target. A potent and selective inhibitor of PDE5 provides vasodilating, relaxing, and diuretic effects, all of which are beneficial in the treatment of various disease states. Research in this  
10 area has led to several classes of inhibitors based on the cGMP basic structure (E. Sybertz et al., *Expert. Opin. Ther. Pat.*, 7, p. 631 (1997)).

The biochemical, physiological, and clinical effects of PDE5 inhibitors therefore suggest  
15 their utility in a variety of disease states in which modulation of smooth muscle, renal, hemostatic, inflammatory, and/or endocrine function is desirable. The compounds of formula (I), therefore, have utility in the treatment of a number of disorders, including stable, unstable, and variant  
20 (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g., postpercutaneous transluminal coronary or carotid angioplasty, or post-bypass surgery graft stenosis), peripheral vascular disease, vascular disorders, such as Raynaud's disease, thrombocytopenia, inflammatory diseases, stroke, bronchitis,  
25 chronic asthma, allergic asthma, allergic rhinitis, glaucoma, osteoporosis, preterm labor, benign prostatic hypertrophy, peptic ulcer, male erectile dysfunction, female sexual dysfunction, and diseases  
30

- 33 -

characterized by disorders of gut motility (e.g., irritable bowel syndrome).

5 An especially important use is the treatment of male erectile dysfunction, which is one form of impotence and is a common medical problem. Impotence can be defined as a lack of power, in the male, to copulate, and can involve an inability to achieve penile erection or ejaculation, or both. The incidence of erectile dysfunction increases with age, with about 50% of men over the age of 40 suffering from some degree of erectile dysfunction.

10 In addition, a further important use is the treatment of female arousal disorder. Female arousal disorders are defined as a recurrent inability to attain or maintain an adequate lubrication/swelling response of sexual excitement until completion of sexual activity. The arousal response consists of vasocongestion in the pelvis, vaginal lubrication, and expansion and swelling of external genitalia.

20 It is envisioned, therefore, that compounds of formula (I) are useful in the treatment of male erectile dysfunction and female arousal disorder. Thus, the present invention concerns the use of compounds of formula (I), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a male animal and arousal disorder in a female animal, including humans.

30 The term "treatment" includes preventing, lowering, stopping, or reversing the progression or

- 34 -

severity of the condition or symptoms being treated. As such, the term "treatment" includes both medical therapeutic and/or prophylactic administration, as appropriate.

5           It also is understood that "a compound of formula (I)," or a physiologically acceptable salt or solvate thereof, can be administered as the neat compound, or as a pharmaceutical composition containing either entity.

10           Although the compounds of the invention are envisioned primarily for the treatment of sexual dysfunction in humans, such as male erectile dysfunction and female arousal disorder, they also can be used for the treatment of other disease states.

15           A further aspect of the present invention, therefore, is providing a compound of formula (I) for use in the treatment of stable, unstable, and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g., post-PTCA or post-bypass graft stenosis), peripheral vascular disease, vascular disorders such as Raynaud's disease, thrombocythemia, inflammatory diseases, prophylaxis of myocardial infarction, prophylaxis of stroke, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, osteoporosis, preterm labor, benign prostatic hypertrophy, male and female erectile dysfunction, or diseases characterized by disorders of gut motility (e.g., IBS).

20

25

30

- 35 -

According to another aspect of the present invention, there is provided the use of a compound of formula (I) for the manufacture of a medicament for the treatment of the above-noted conditions and disorders.

In a further aspect, the present invention provides a method of treating the above-noted conditions and disorders in a human or nonhuman animal body which comprises administering to said body a therapeutically effective amount of a compound of formula (I).

Compounds of the invention can be administered by any suitable route, for example by oral, buccal, inhalation, sublingual, rectal, vaginal, transurethral, nasal, topical, percutaneous, i.e., transdermal, or parenteral (including intravenous, intramuscular, subcutaneous, and intracoronary) administration. Parenteral administration can be accomplished using a needle and syringe, or using a high pressure technique, like POWDERJECT™.

Oral administration of a compound of the invention is the preferred route. Oral administration is the most convenient and avoids the disadvantages associated with other routes of administration. For patients suffering from a swallowing disorder or from impairment of drug absorption after oral administration, the drug can be administered parenterally, e.g., sublingually or buccally.

Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effec-

- 36 -

tive amount" means an amount effective to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in  
5 light of the detailed disclosure provided herein.

A "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired effect. Toxicity and therapeutic efficacy of such compounds can be determined  
10 by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in  
15 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD<sub>50</sub> and ED<sub>50</sub>. Compounds which exhibit high therapeutic indices are preferred. The data obtained from such  
20 data can be used in formulating a range of dosage for use in humans. The dosage of such compounds preferably lies within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage can vary within this range  
25 depending upon the dosage form employed, and the route of administration utilized.

The exact formulation, route of administration, and dosage can be chosen by the individual physician in view of the patient's condition.  
30 Dosage amount and interval can be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the therapeutic effects.

- 37 -

The amount of composition administered is dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician.

Specifically, for administration to a human in the curative or prophylactic treatment of the conditions and disorders identified above, oral dosages of a compound of formula (I) generally are about 0.5 to about 1000 mg daily for an average adult patient (70 kg). Thus, for a typical adult patient, individual tablets or capsules contain 0.2 to 500 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier, for administration in single or multiple doses, once or several times per day. Dosages for intravenous, buccal, or sublingual administration typically are 0.1 to 500 mg per single dose as required. In practice, the physician determines the actual dosing regimen which is most suitable for an individual patient, and the dosage varies with the age, weight, and response of the particular patient. The above dosages are exemplary of the average case, but there can be individual instances in which higher or lower dosages are merited, and such are within the scope of this invention.

For human use, a compound of the formula (I) can be administered alone, but generally is administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. Pharmaceutical compositions for use in accordance with the present invention thus can be

- 38 -

formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries that facilitate processing of compounds of formula (I) into preparations  
5 which can be used pharmaceutically.

These pharmaceutical compositions can be manufactured in a conventional manner, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Proper  
10 formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of a compound of the present invention is administered orally, the composition typically is in the form of a tablet, capsule, powder, solution, or  
15 elixir. When administered in tablet form, the composition can additionally contain a solid carrier, such as a gelatin or an adjuvant. The tablet, capsule, and powder contain about 5% to about 95% compound of the present invention, and preferably from  
20 about 25% to about 90% compound of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, or oils of animal or plant origin can be added. The liquid form of the composition can further contain  
25 physiological saline solution, dextrose or other saccharide solutions, or glycols. When administered in liquid form, the composition contains about 0.5% to about 90% by weight of a compound of the present invention, and preferably about 1% to about 50% of a  
30 compound of the present invention.

When a therapeutically effective amount of a compound of the present invention is administered

- 39 -

by intravenous, cutaneous, or subcutaneous injection, the composition is in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred composition for intravenous, cutaneous, or subcutaneous injection typically contains, in addition to a compound of the present invention, an isotonic vehicle.

For oral administration, the compounds can be formulated readily by combining a compound of formula (I) with pharmaceutically acceptable carriers well known in the art. Such carriers enable the present compounds to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding a compound of formula (I) with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers and cellulose preparations. If desired, disintegrating agents can be added.

For administration by inhalation, compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered



- 40 -

amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

5           The compounds can be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, e.g., in ampules or in multidose containers, 10 with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing, and/or dispersing agents.

15           Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. 20 Suitable lipophilic solvents or vehicles include fatty oils or synthetic fatty acid esters. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension. Optionally, the suspension also can contain suitable 25 stabilizers or agents that increase the solubility of the compounds and allow for the preparation of highly concentrated solutions. Alternatively, a present composition can be in powder form for constitution with a suitable vehicle, e.g., sterile 30 pyrogen-free water, before use.

Compounds of the present invention also can be formulated in rectal compositions, such as suppositories or retention enemas, e.g., containing

- 41 -

conventional suppository bases. In addition to the formulations described previously, the compounds also can be formulated as a depot preparation. Such long-acting formulations can be administered by  
5 implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion  
10 exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Many of the compounds of the present invention can be provided as salts with pharmaceutically compatible counterions. Such pharmaceutically  
15 acceptable base addition salts are those salts that retain the biological effectiveness and properties of the free acids, and that are obtained by reaction with suitable inorganic or organic bases.

In particular, a compound of formula (I)  
20 can be administered orally, buccally, or sublingually in the form of tablets containing excipients, such as starch or lactose, or in capsules or ovules, either alone or in admixture with excipients, or in the form of elixirs or suspensions containing  
25 flavoring or coloring agents. Such liquid preparations can be prepared with pharmaceutically acceptable additives, such as suspending agents. A compound also can be injected parenterally, for example, intravenously, intramuscularly, subcutaneously, or intracoronarily. For parenteral admin-  
30 istration, the compound is best used in the form of a sterile aqueous solution which can contain other substances, for example, salts, or monosaccharides,

- 42 -

such as mannitol or glucose, to make the solution isotonic with blood.

For veterinary use, a compound of formula (I) or a nontoxic salt thereof, is administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal.

Thus, the invention provides in a further aspect a pharmaceutical composition comprising a compound of the formula (I), together with a pharmaceutically acceptable diluent or carrier therefor. There is further provided by the present invention a process of preparing a pharmaceutical composition comprising a compound of formula (I), which process comprises mixing a compound of formula (I), together with a pharmaceutically acceptable diluent or carrier therefor.


In a particular embodiment, the invention includes a pharmaceutical composition for the curative or prophylactic treatment of erectile dysfunction in a male animal, or arousal disorder in a female animal, including humans, comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

Compounds of formula (I) can be prepared by any suitable method known in the art, or by the following processes which form part of the present invention. In the methods below,  $R^0$ ,  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$ , as well as  $R^a$  through  $R^n$ , Q, B, C, D, E, X, and Y, are defined as in structural formula (I) above.

- 43 -

In particular, compounds of structural formula (I) can be prepared according to the following synthetic scheme.

5 Daugan U.S. Patent No. 5,859,006, incorporated herein by reference, discloses preparation of a compound of structural formula (III):



(III)



15

The compounds of structural formula (I) can be prepared from compounds similar to the compound of structural formula (III) using appropriate  $R^2$  and  $R^4$  substituents.


20

The following illustrates a general method of synthesizing a compound of structural formula (I).

- 44 -



- 45 -



5

10

15


Known compound (IV) (see Daugan U.S. Patent No. 5,859,006) is hydrolyzed to carboxylic acid (V), which then is treated with triphosgene followed by treatment with a desired amine to pro-

20


- 46 -

vide amide (VI). Amide (VI) then is treated with a carboxylic acid in the presence of an activating group, or other carboxylic acid derivative (e.g., acid chloride), having the desired  $-(X)_n-Y$  residue and a base to provide bisamide (VII). Bisamide (VII) is a compound of structural formula (I).

The following illustrates a synthesis of compounds (V)-(VII) from compound (IV), wherein  $R^0$  is hydrogen. The synthesis of compound (IV) can be found in Daugan U.S. Patent No. 5,859,006.




(IVa) (IV, wherein  $R^0=H$ )



(Va) (V, wherein  $R^0=H$ )

- 47 -



5

10

15

20



- 48 -

**Preparation of (1R,3R)-1-benzo[1,3]dioxol-5-yl-2,3,4,9-tetrahydro-1H- $\beta$ -carboline-3-carboxylic acid hydrochloride (Va)**

5                   Lithium hydroxide (0.32 g, 13.2 mmol) was added to a stirred mixture of (1R,3R)-1-benzo[1,3]-dioxol-5-yl-2,3,4,9-tetrahydro-1H- $\beta$ -carboline-3-carboxylic acid methyl ester (IVa) (2.5 g, 6.5 mmol)  
10 in 10 mL of anhydrous tetrahydrofuran at room temperature. Methanol (1 mL) was added to help dissolution of the solids. After stirring overnight at room temperature, the reaction mixture was partitioned between ethyl acetate (10 mL) and deionized  
15 water (25 mL). The layers were separated, and the pH of the aqueous layer was adjusted to 2.3 with 1 M HCl. After 1 hour, the resulting slurry was filtered, then the filter cake was washed with a quantity of deionized water sufficient to remove the  
20 yellow color. The resulting solid was dried under vacuum to give 1.38 g (57%) of compound (Va) as an off-white solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ : 10.70 (s, 1H), 7.52 (d, J=7 Hz, 1H), 7.28 (d, J=8 Hz, 1H), 7.13-7.01 (m, 5H), 6.07  
25 (s, 2H), 5.78 (s, 1H), 4.22 (dd, J=5, 11 Hz, 1H), 3.38 (dd, J=5, 15 Hz, 1H), 3.21 (dd, J=2, 15 Hz, 1H); MS ES+m/e 337 (p+1), ES-m/e 335 (p-1); IR (KBr, cm<sup>-1</sup>) 3621, 3450, 1758.

30                   **Preparation of (1R,3R)-1-benzo[1,3]dioxol-5-yl-2-(3-phenyl-acryloyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline-3-carboxylic acid dimethylamide (VIa)**

35                   Triethylamine (1.5 mL, 10.7 mmol) was added to a stirred mixture of compound (Va) (2 g,

- 49 -

5.4 mmol) and 60 mL of anhydrous tetrahydrofuran at room temperature. Triphosgene (1.6 g, 5.4 mmol) was added to the slurry, then stirring was continued for 2 hours at room temperature. Dimethylamine was condensed into a dropping funnel (about 1 mL), and was added directly to the stirred slurry. After 1 hour, the reaction mixture was filtered and the volatiles were removed from the filtrate under vacuum. The resulting foam was subjected to flash chromatography (silica, EtOAc) to give a 1.65 g (86%) of compound (VIa) as a yellow foam.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.38 (s, 1H), 7.44 (d, J=6 Hz, 1H), 7.21 (d, J=7 Hz, 1H), 7.03-6.88 (m, 3H), 6.83-6.79 (m, 2H), 5.99 (s, 2H), 5.13 (d, J=7 Hz, 1H), 4.04 (m, 1H), 3.12 (s, 3H), 2.86 (s, 3H), 2.8-2.76 (m, 2H); MS ES+m/e 364 (p+1), ES-m/e 362 (p-1); IR (KBr, cm<sup>-1</sup>) 3463, 1642.

**Preparation of (1R,3R)-1-benzo[1,3]dioxol-5-yl-2-(3-phenylacryloyl)-2,3,4,9-tetrahydro-1H-β-carboline-3-carboxylic acid dimethylamide (VIIa)**


2-Chloro-3,5-dimethoxy-2,4,6-triazine (0.5 g, 2.8 mmol) was added to a mixture of trans-cinnamic acid in 15 mL of anhydrous tetrahydrofuran. N-Methylmorpholine (0.32 mL, 2.9 mmol) was added and the resulting solution was stirred at room temperature for 1.75 hours. Compound (VIa) was added, then the reaction was stirred for 4 hours at room temperature. The reaction mixture was partitioned between ethyl acetate (25 mL) and 1 M HCl (25 mL). The layers were separated, and the organic layer was washed with 25 mL of saturated sodium bicarbonate

- 50 -


solution followed by 25 mL of brine. The EtOAc was dried over  $\text{MgSO}_4$ , filtered, and removed under vacuum to give the crude product as a yellow foam. The product was purified by flash chromatography (50% hexane, 50%  $\text{CH}_2\text{Cl}_2$  to  $\text{CH}_2\text{Cl}_2$  to EtOAc) to yield 0.613 g (42%) of compound (VIIa) as a yellow foam: mp 175-178°C.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$ : 10.82, 10.75 (overlapping br s, 1H), 7.89-7.21 (m, 9H), 7.11-6.67 (m, 5H), 6.65 (br s, 1H), 5.92 (s, 2H), 5.70-5.63 (m, 1H), 2.9 (dd,  $J=6, 15$  Hz, 1H), 2.75 (br s, 1H), 2.58 (br s, 2H), 2.51-2.48 (m, 1H), 2.36 (br s, 2H), 2.18 (br s, 1H); MS ES-m/e 492 (p-1).


The following scheme illustrates the synthesis of a compound of structural formula (I) from ester (IV), and illustrates an alternative method of synthesizing a bisamide (VII).



- 51 -



(VIII)



Example 1

The above scheme illustrates the synthesis of a compound of structural formula (II), i.e., compound (VIII), directly from compound (IVa) by reaction with a carboxylic acid having the desired (X)<sub>n</sub>-Y residue, in the presence of dicyclohexylcarbodiimide and triethylamine. Compound (VIII) then can be converted to another compound of structural formula (I), i.e., Example 1, by a reaction using trimethyl aluminum and dimethylamine hydrochloride. The

- 52 -

reaction between compound (IVa) and cinnamic acid is set forth in Bombrun U.S. Patent No. 6,117,881, incorporated in its entirety herein by reference.

It should be understood that protecting groups can be utilized in accordance with general principles of synthetic organic chemistry to provide compounds of structural formula (I). Protecting group-forming reagents, like benzyl chloroformate and trichloroethyl chloroformate, are well known to persons skilled in the art, for example, see T.W. Greene et al., "Protective Groups in Organic Synthesis, Third Edition," John Wiley and Sons, Inc., NY, NY (1999). These protecting groups are removed when necessary by appropriate basic, acidic, or hydro-

5 groups can be utilized in accordance with general  
principles of synthetic organic chemistry to provide  
compounds of structural formula (I). Protecting  
group-forming reagents, like benzyl chloroformate  
and trichloroethyl chloroformate, are well known to  
persons skilled in the art, for example, see T.W.  
Greene et al., "Protective Groups in Organic Synthe-  
sis, Third Edition," John Wiley and Sons, Inc., NY,  
NY (1999). These protecting groups are removed when  
necessary by appropriate basic, acidic, or hydro-  
genolytic conditions known to persons skilled in the  
art. Accordingly, compounds of structural formula  
(I) not specifically exemplified herein can be pre-  
pared by persons skilled in the art.

10  
15

In addition, compounds of formula (I) can be converted to other compounds of formula (I). Thus, for example, a particular R substituent can be interconverted to prepare another suitably substituted compound of formula (I). Examples of appropriate interconversions include, but are not limited to, OR<sup>a</sup> to hydroxy by suitable means (e.g., using a agent such as BBr<sub>3</sub>, or a palladium catalyst, like palladium-on-carbon, and hydrogen), or amino to substituted amino, such as acylamino or sulphonyl-amino, using standard acylating or sulfonylating conditions.

20  
25  
30

Compounds of formula (I) can be prepared by the method above as individual stereoisomers or as a racemic mixture. Individual stereoisomers of

- 53 -


the compounds of the invention can be prepared from racemates by resolution using methods known in the art for the separation of racemic mixtures into their constituent stereoisomers, for example, using  
5 HPLC on a chiral column, such as Hypersil naphthyl urea, or using separation of salts of stereoisomers. Compounds of the invention can be isolated in association with solvent molecules by crystallization from, or evaporation of, an appropriate solvent.  
10

The pharmaceutically acceptable acid addition salts of the compounds of formula (I) that contain a basic center can be prepared in a conventional manner. For example, a solution of the free base  
15 can be treated with a suitable acid, either neat or in a suitable solution, and the resulting salt isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts can be obtained in an analogous manner by treating a solution  
20 of a compound of formula (I) with a suitable base. Both types of salt can be formed or interconverted using ion-exchange resin techniques. Thus, according to a further aspect of the invention, a method  
25 for preparing a compound of formula (I) or a salt or solvate (e.g., hydrate) is provided, followed by (i) salt formation, or (ii) solvate (e.g., hydrate) formation.


The following additional abbreviations are  
30 used hereafter in the accompanying examples: rt (room temperature), min (minute), h (hour), g (gram), mmol (millimole), m.p. (melting point), LiOH (lithium hydroxide), eq (equivalents), L (liter), mL

- 54 -


(milliliter),  $\mu\text{L}$  (microliter), DMSO (dimethyl sulfoxide),  $\text{Et}_3\text{N}$  (triethylamine),  $\text{MeNH}_2$  (methylamine), THF (tetrahydrofuran),  $\text{Me}_2\text{NH}$  (dimethylamine), DCC (1,3-dicyclohexylcarbodiimide),  $\text{AlMe}_3$  (trimethylaluminum), Me (methyl), EtOAc (ethyl acetate),  $\text{CHCl}_3$  (chloroform), and  $\text{Na}_2\text{SO}_4$  (sodium sulfate).

**Example 2**

20 Example 2 was prepared from Compound (IVa) by the following synthetic sequence.




- 55 -

**Example 3**

5     (-) - (1R,3R) - 1-Benzo[1,3]dioxol-5-yl-1,3,4,9-tetra-  
hydro-beta-carboline-2,3-dicarboxylic acid 3-methyl  
ester 2-(4-nitrophenyl)ester

10

15




20

Example 3 was prepared from Compound (IVa) as follows. Also see F. Pinnen et al., *J. Chem. Soc., Perkin Trans. I*, 12, p. 1611 (1994).



- 56 -




4-Nitrophenyl chloroformate was added dropwise to a suspension of Compound (IVa) (2.0 g, 52 mmol) and Et<sub>3</sub>N (1.8 mL, 13 mmol) in THF (50 mL) at 0°C under a nitrogen blanket. The resulting mixture was slowly warmed to room temperature and stirred for 18 hours, then diluted with EtOAc (200 mL). The resulting solution was washed with brine (100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography, eluting with EtOAc/CHCl<sub>3</sub> (1:19) to provide Example 3 as a yellow powder (2.08 g, 78%): mp 175-187°C; TLC R<sub>f</sub> (9:1 CHCl<sub>3</sub>/EtOAc)=0.84.


<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ: 10.90 (s, 1H), 8.33 (d, J=9.0 Hz, 2H), 7.58-7.56 (m, 3H), 7.30 (d, J=7.7 Hz, 1H), 7.07 (dt, J=6.5, 21.8 Hz, 2H), 6.88 (bs, 1H), 6.72 (s, 1H), 6.58 (bs, 1H), 6.44 (bs, 1H), 6.01 (s, 2H), 5.70 (bs, 1H), 5.45 (bs, 1H), 3.40 (d, J=13.7 Hz, 1H), 3.28-3.21 (m, 1H); API MS m/z 516 [C<sub>27</sub>H<sub>21</sub>N<sub>3</sub>O<sub>8</sub>+H]<sup>+</sup>; [α]<sub>D</sub><sup>25°C</sup>=-121.7° (c=1.0, DMSO). Anal. Calcd. for C<sub>27</sub>H<sub>21</sub>N<sub>3</sub>O<sub>8</sub>: C, 62.91; H, 4.11; N, 8.15. Found: C, 62.54; H, 3.91; N, 8.02.

The following Examples 4-12 were prepared in a manner similar to Examples 1-3.

- 57 -




Example 4



Example 5

- 58 -

5




10

Example 6

15

20




25

Example 7

30


- 59 -

### Preparation of Example 8



### Example 8

Example 8 was prepared by the following synthetic sequence



20

- 60 -

**Preparation of (1R)-(+)-1-benzo[1,3]dioxol-5-yl-  
2,3,4,9-tetrahydro-1H- $\beta$ -carboline-3-carboxylic  
acid propylamide (Compound (IX))**

5                   A suspension of compound (Va) (2.0 g, 6.0 mmol) and triethylamine (2 mL) in tetrahydrofuran (50 mL) was immersed in a sonicator until all solids dissolved. Triphosgene (1.6 g, 5.4 mmol) was added,  
10                   and the resulting suspension was stirred for 6 hours without cooling. A colorless solid was removed by filtration, then the filtrate was concentrated to a foam. The foam was dissolved in chloroform (50 mL) and n-propylamine (2 mL) was added. The reaction  
15                   was stirred for 18 hours without cooling, the solvent was evaporated, and the residue purified by chromatography (silica gel, 75% ethyl acetate: 25% hexanes) to give 1.4 g (61%) of compound (IX) as a solid.

20                   <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ : 10.71 (s, 1H), 7.84 (t, J=8 Hz, 1H), 7.45-6.59 (m, 7H), 6.0 (s, 2H), 5.15 (s, 1H), 3.45 (m, 1H), 3.04 (q, J=8 Hz, 2H), 2.8 (dd, J=5, 13 Hz, 1H), 1.4 (m, 2H), 0.84 (t, J=8 Hz, 3H); MS ES+m/e 378.2 (p+1), ES-m/e 376.2 (p-1); Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>; C, 70.01; H, 6.14; N, 11.13. Found:  
25                   C, 70.30; H, 6.23; N, 11.27.

30                   **Preparation of (1R)-(+)-2-acetyl-1-benzo[1,3]dioxol-5-yl-2,3,4,9-tetrahydro-1H- $\beta$ -carboline-3-carboxylic acid propylamide (Example 8)**


                  To a solution of compound (IX) (1.4 g, 3.7 mmol) and triethylamine (2 mL) in methylene chloride  
35                   (25 mL) was added acetyl chloride (0.36 mL, 5.0 mmol), then the mixture was stirred for 18 hours at

- 61 -

room temperature. The solution was washed once with water, once with saturated sodium chloride solution, dried (sodium sulfate), filtered, and concentrated *in vacuo* to provide a solid that was recrystallized (ethyl acetate) to give 310 mg (20%) of Example 8 as a solid: mp 163-165°C.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.88 (s, 1H), 7.95 (t, J=5 Hz, 0.6H (rotamer)), 7.7 (t, J=5 Hz, 0.4H (rotamer)), 7.4-6.7 (m, 7H), 5.85-6.2 (m, 3H), 5.25 (br s, 0.6H (rotamer)), 4.95 (br s, 0.4 H (rotamer)), 3.35 (dd, J=6, 13 Hz, 2H), 2.85 (m, 2H), 2.0 (s, 1.8H (rotamer)), 1.9 (s, 1.2H (rotamer)), 1.2 (m, 2H), 0.62 (m, 3H); MS ES+m/e 420.2 (p+1), ES-m/e 418.3 (p-1); Anal. Calcd. for C<sub>24</sub>H<sub>25</sub>N<sub>3</sub>O<sub>4</sub>: C, 68.71; H, 6.01; N, 10.02. Found: C, 68.57; H, 5.69; N, 10.02.


#### Preparation of Example 9



Example 9

- 62 -

Example 9 was prepared by the following synthetic sequence:



5

**Preparation of (1R) - (+) - (1-benzo[1,3]dioxol-5-yl-2,3,4,9-tetrahydro-1H- $\beta$ -carbolin-3-yl) - morpholine-4-yl-methanone (X)**

10

A suspension of compound (Va) (350 mg, 1.04 mmol) and triethylamine (0.5 mL) in tetrahydrofuran (10 mL) was sonicated until all solids were dissolved. Triphosgene (300 mg, 1.0 mmol) was added and the resulting suspension was stirred for 2 hours. A colorless solid was removed by filtration and the filtrate was concentrated *in vacuo* to provide a foam. The foam was dissolved in methylene chloride (15 mL), treated with morpholine (0.5 mL), and the resulting solution stirred for 18 hours. The solvent was evaporated *in vacuo*, and the crude material was purified by chromatography (silica gel,

- 63 -

ethyl acetate) to give 280 mg (66%) of compound (X) as a solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.35 (s, 1H), 7.45-6.77 (m, 7H), 6.02 (s, 2H), 5.15 (s, 1H), 4.06 (dd, J=4, 13 Hz), 3.7-3.45 (m, 8H), 2.85 (dd, J=4, 13 Hz, 2H), 2.1 (t, J=10 Hz, 1H); MS ES+m/e 406.0 (p+1), ES-m/e 404.2 (p-1).

10 **Preparation of (1R)-(+)-1-[benzo[1,3]dioxol-5-yl-3-(morpholine-4-carbonyl)-1,3,4,9-tetrahydro-β-carbolin-2-yl]-ethanone (Example 11)**


15 To a solution of compound (X) (1.7 g, 4.2 mmol) and triethylamine (2 mL) in methylene chloride (50 mL) was added acetyl chloride (3.29 mg, 4.2 mmol, 0.3 mL). The reaction was stirred for 18 hours without cooling. An additional quantity of acetyl chloride (0.1 mL) was added, and stirring was  
20 continued for 30 minutes. The reaction was concentrated and purified by chromatography (silica gel, 75% ethyl acetate: 25% hexanes, then ethyl acetate) to give 800 mg (42%) of Example 9 as a colorless solid.

25 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.8 (s, 1H), 7.55-6.8 (m, 7H), 6.75 (br s, 0.6H (rotamer)), 6.62 (br s, 0.4H (rotamer)), 5.97 (d, J=4 Hz, 2H), 5.6 (br s, 0.6H (rotamer)), 5.75 (br s, 0.4H (rotamer)), 3.54-2.95 (m, 9H), 2.85 (dd, J=5, 13 Hz, 1H), 2.3 (s, 3H); MS  
30 ES+m/e 448.1 (p+1), ES-m/e 446.2 (p-1); Anal. Calcd. for C<sub>25</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>: C, 67.10; H, 5.63; N, 9.39. Found: C, 67.00; H, 5.69; N, 9.28.



- 64 -

5




10

Example 10

Preparation of Example 11

15

20




25

Example 11


Example 11 was prepared using the  
following synthetic sequence:

30

- 65 -

trans-cinnamic acid

N-methylmorpholine

 $\longrightarrow$  Example 11

Preparation of (1R,3R)-(+)-1-benzo[1,3]dioxol-5-yl-  
2,3,4,9-tetrahydro-1H-β-carboline-3-carboxylic acid  
butylamide (XI)

5

To a solution of compound (Va) (2.17 g, 6  
mmol) in chloroform (50 mL) was added n-butylamine  
10 (585 mg, 6 mmol). The reaction was stirred for 18  
hours at room temperature. The solvent was con-  
centrated in vacuo, then the residue was purified by

- 66 -

chromatography (silica gel, 50% ethyl acetate: 70% hexanes) to give 1.7 g (72%) of compound (XI) as a colorless foam.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.28 (s, 1H), 7.6 (t, J=6 Hz, 1H), 7.45-6.85 (m, 8H), 6.0 (s, 2H), 5.08 (d, J=7 Hz, 1H), 3.55 (m, 1H), 3.14 (q, J=6 Hz, 2H), 2.97 (dd, J=3, 16 Hz, 1H), 2.7 (dt, J=3, 16 Hz), 1.47 (m, 2H), 1.3 (m, 2H), 0.82 (t, J=6 Hz, 3H); MS ES+m/e 392.3 (p+1), ES-m/e 390.4 (p-1); Anal. Calcd. for C<sub>23</sub>H<sub>25</sub>N<sub>3</sub>O<sub>3</sub>: C, 70.57; H, 6.43; N, 10.73. Found: C, 70.34; H, 6.11; N, 10.53.

**Preparation of (1R,3R)-(+)-1-benzo[1,3]dioxol-5-yl-2-(3-phenylacryloyl)-2,3,4,9-tetrahydro-1H-β-carboline-3-carboxylic acid butylamide (Example 11)**

A mixture of trans-cinnamic acid (376 mg, 2.5 mmol), 2-chloro-3,5-dimethoxy-2,4,6-triazine (500 mg, 2.8 mmol), N-methylmorpholine (0.32 mL, 2.9 mmol), and tetrahydrofuran (20 mL) was stirred without cooling for 1.5 hours. Compound (XI) (1.0 g, 2.5 mmol) was added, then the reaction was stirred for an additional 18 hours. The solvent was evaporated, and the residue was dissolved in ethyl acetate, washed once with 1N hydrochloric acid, once with saturated sodium bicarbonate solution, dried (sodium sulfate), filtered, and concentrated in vacuo. Chromatography (silica gel, 30% ethyl acetate: 70% hexanes) of the residue provided 800 mg (61%) of Example 11 as a solid: mp 114-117°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.8 (s, 1H), 6.8-7.7 (m, 14H), 5.95 (s, 2H), 5.3 (m, 1H), 3.0 (m, 1H), 2.78 (m, 1H), 1.15 (m, 6H), 0.8 (m, 3H); MS ES+m/e 522.02

- 67 -


(p+1), ES-m/e 520.13 (p-1), IR (KBr,  $\text{cm}^{-1}$ ) 1674, 1644, 1503, 1487; Anal. Calcd. for  $\text{C}_{32}\text{H}_{31}\text{N}_3\text{O}_4$ : C, 73.68; H, 5.99; N, 8.06. Found: C, 73.33; H, 5.96; N, 8.25.

5

Preparation of Example 12

10

15



Example 12


20

Example 12 was prepared using the following synthetic sequence:

25


30

- 68 -



$\xrightarrow{\hspace{1.5cm}}$  Example 12

trans-cinnamic acid,  
N-methylmorpholine



Preparation of (1R,3R)-1-benzo[1,3]dioxol-5-yl-  
2-(3-phenylacryloyl)-2,3,4,9-tetrahydro-1H-β-  
carboline-3-carboxylic acid methylamide (XII)

Triethylamine (2 mL, 14 mm, 2.4 eq) was added to a stirred mixture of compound (Va) (2 g, 5.2 mmol) and anhydrous tetrahydrofuran (25 mL) at room temperature. The resulting mixture was sonicated until complete dissolution was achieved.

- 69 -

Triphosgene (1.6 g, 5.4 mmol) was added to the slurry, and stirring was continued for 5 hours at room temperature. The solids were removed by filtration and the filtrate concentrated *in vacuo*.

5 The residue was dissolved in chloroform, and a solution of 2 M methylamine in tetrahydrofuran (10 mL, 20 mmol) was added. After 1.5 hours, the reaction mixture was diluted with water and shaken. The organic layer was dried (sodium sulfate),  
10 filtered, and concentrated *in vacuo*. The residue (compound (XII)) was used as is.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 10.38 (s, 1H), 7.44 (d, J=6 Hz, 1H), 7.21 (d, J=7 Hz, 1H), 7.03-6.88 (m, 3H), 6.83-6.79 (m, 2H), 5.99 (s, 2H), 5.10 (m, 1H), 3.5-3.6  
15 (m, 1H), 3.30 (s, 3H), 3.00 (dd, J=1.5, 11.4 Hz, 1H), 2.72 (dd, J=2.5, 11.3 Hz, 1H), 2.64 (s, 3H), 2.49 (s, 3H); MS ES+m/e 350.2 (p+1), ES-348.2 (p-1).


20 **Preparation of (1R,3R)-{2-[(2E)-3-(phenyl)prop-2-enoyl]-1-(2H-benzo[d]1,3-dioxolan-5-yl)-(1,2,3,4-tetrahydrobetacarbolin-3-yl)}-N-methylcarboxamide (Example 12)**

25 2-Chloro-3,5-dimethoxy-2,4,6-triazine (0.5 g, 2.8 mmol) was added to a solution of trans-cinnamic acid (0.376 g, 2.5 mmol) in anhydrous tetrahydrofuran (20 mL). N-Methylmorpholine (0.32 mL, 2.9 mmol) was added, then the resulting solution  
30 was stirred at room temperature for 1.5 hours. A solution of compound (XII) (0.873 g, 2.50 mmol) in anhydrous tetrahydrofuran (5 mL) was added, and the reaction was stirred for 2 hours at room temperature. The reaction mixture was partitioned between  
35 ethyl acetate and 1M hydrochloric acid. The layers

- 70 -

were separated, and the organic layer was washed once with saturated sodium bicarbonate solution and once with saturated sodium chloride solution. The organic layer was dried (sodium sulfate), filtered, and concentrated *in vacuo* to give the crude product as a yellow foam. The product was purified by flash chromatography (silica gel, 50% hexane/50% ethyl acetate to ethyl acetate) to provide 0.43 g (35%) of Example 12 as a yellow foam. MS FAB exact mass calculated for  $C_{29}H_{23}N_4O_4$ :  $m/z=479.1845$ . Found: 479.1845.

#### Preparation of Example 13



Example 13

Example 13 was prepared in an identical manner to Example 12 and substituting p-chloro *trans* cinnamic acid for *trans* cinnamic acid.


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ : 2.18 (br s, 1.5H), 2.35 (br s, 1H), 2.50 (m, 1H), 2.51 (br s, 2H), 2.73 (br s, 1H),

- 71 -

2.77 (dd, J=5.8 Hz, 15 Hz, 1H), 3.24-3.40 (m, 0.5H),  
5.75 (br s, 1H), 5.95 (s, 2H), 6.68 (br s, 1H),  
6.78-7.10 (m, 5H), 7.26-7.37 (m, 1H), 7.46-7.50 (m,  
7H), 10.75 (br s, 0.5H), 10.88 (br s, 0.5H); MS  
5 ES+m/e 528.2 (p+1), MS ES-m/e 526.2 (p-1).

The following intermediates have been prepared, and can be used to provide a composition of structural formula (I) by the methods set forth above. In particular, an intermediate is reacted  
10 with a carboxylic acid, or a carboxylic acid derivative, having the desired (X)<sub>n</sub>-Y residue. As demonstrated hereafter, the following intermediates also are potent and selective inhibitors of PDE5, and, like compounds of structural formula (I), can be  
15 used in a variety of therapeutic areas wherein such inhibition is considered beneficial.

20




Intermediate 1a



- 72 -

5




10

Intermediate 1b

15


20



Intermediate 2

- 73 -


5



Intermediate 3


10

15



20


Intermediate 4



Intermediate 5

- 74 -


5



10

Intermediate 6


15



20

Intermediate 7a

25




30

Intermediate 7b

- 75 -


5



10

Intermediate 8


15



20

Intermediate 9a

25




30

Intermediate 9b

- 76 -


5



10

Intermediate 10


15



20

Intermediate 11

25




30

Intermediate 12

- 77 -

5

10




Intermediate 13a

15

20


25




Intermediate 13b

30

- 78 -




Intermediate 14



Intermediate 15a

- 79 -

5




10

Intermediate 15b

15

20



25


Intermediate 16a

30



- 80 -


5



10

Intermediate 16b

15




20

25


Intermediate 17a

30

- 81 -




Intermediate 17b



Intermediate 18

- 82 -

5




10

Intermediate 19

15

20




25

Intermediate 20a

30

- 83 -

5




10

Intermediate 20b

15

20




25

Intermediate 21

30

- 84 -

5




10

Intermediate 22

15

20




25


Intermediate 23

30

- 85 -




Intermediate 24a




Intermediate 24b

30

- 86 -




Intermediate 25



Intermediate 26a

- 87 -

5




10

Intermediate 26b

15

20




25

Intermediate 27


30



- 88 -



Intermediate 28




Intermediate 29a

25

30

- 89 -


5



10

Intermediate 29b


15



20

Intermediate 30a

25




30

Intermediate 30b

- 90 -


5



10

Intermediate 31

15



20

25


Intermediate 32

30

- 91 -

5

10




Intermediate 33

15

20

25




Intermediate 34

30

- 92 -

5


10



Intermediate 35


15

20




Intermediate 36

25



Intermediate 37


20



Intermediate 38

30

- 94 -



Intermediate 39

Compounds of the present invention can be formulated into tablets for oral administration. For example, a compound of formula (I) can be formed into a dispersion with a polymeric carrier by the coprecipitation method set forth in WO 96/38131, incorporated herein by reference. The coprecipitated dispersion then can be blended with excipients, then pressed into tablets, which optionally are film-coated.

The compounds of structural formula (I) were tested for an ability to inhibit PDE5. The ability of a compound to inhibit PDE5 activity is related to the  $IC_{50}$  value for the compound, i.e., the concentration of inhibitor required for 50% inhibition of enzyme activity. The  $IC_{50}$  value for compounds of structural formula (I) were determined using recombinant human PDE5.

The compounds of the present invention typically exhibit an  $IC_{50}$  value against recombinant human PDE5 of less than about 50  $\mu M$ , and preferably

- 95 -

less than about 25  $\mu\text{M}$ , and more preferably less than about 15  $\mu\text{M}$ . The compounds of the present invention typically exhibit an  $\text{IC}_{50}$  value against recombinant human PDE5 of less than about 1  $\mu\text{M}$ , and often less than about 0.5  $\mu\text{M}$ . To achieve the full advantage of the present invention, a present PDE5 inhibitor has an  $\text{IC}_{50}$  of about 0.1 nM to about 15  $\mu\text{M}$ .

The production of recombinant human PDEs and the  $\text{IC}_{50}$  determinations can be accomplished by well-known methods in the art. Exemplary methods are described as follows:

#### EXPRESSION OF HUMAN PDEs

##### Expression in *Saccharomyces cerevisiae* (Yeast)

Recombinant production of human PDE1B, PDE2, PDE4A, PDE4B, PDE4C, PDE4D, PDE5, and PDE7 was carried out similarly to that described in Example 7 of U.S. Patent No. 5,702,936, incorporated herein by reference, except that the yeast transformation vector employed, which is derived from the basic ADH2 plasmid described in Price et al., *Methods in Enzymology*, 185, pp. 308-318 (1990), incorporated yeast ADH2 promoter and terminator sequences and the *Saccharomyces cerevisiae* host was the protease-deficient strain BJ2-54 deposited on August 31, 1998 with the American Type Culture Collection, Manassas, Virginia, under accession number ATCC 74465. Transformed host cells were grown in 2X SC-leu medium, pH 6.2, with trace metals, and vitamins. After 24 hours, YEP medium-containing glycerol was added to a final concentration of 2X YET/3% glycerol. Approxi-



- 96 -

mately 24 hr later, cells were harvested, washed, and stored at -70°C.

#### HUMAN PHOSPHODIESTERASE PREPARATIONS

5

##### Phosphodiesterase Activity Determinations

Phosphodiesterase activity of the preparations was determined as follows. PDE assays utilizing a charcoal separation technique were performed essentially as described in Loughney et al. (1996). In this assay, PDE activity converts [32P]cAMP or [32P]cGMP to the corresponding [32P]5'-AMP or [32P]5'-GMP in proportion to the amount of PDE activity present. The [32P]5'-AMP or [32P]5'-GMP then was quantitatively converted to free [32P]phosphate and unlabeled adenosine or guanosine by the action of snake venom 5'-nucleotidase. Hence, the amount of [32P]phosphate liberated is proportional to enzyme activity. The assay was performed at 30°C in a 100  $\mu$ L reaction mixture containing (final concentrations) 40 mM Tris HCl (pH 8.0), 1  $\mu$ M ZnSO<sub>4</sub>, 5 mM MgCl<sub>2</sub>, and 0.1 mg/mL bovine serum albumin (BSA). PDE enzyme was present in quantities that yield <30% total hydrolysis of substrate (linear assay conditions). The assay was initiated by addition of substrate (1 mM [32P]cAMP or cGMP), and the mixture was incubated for 12 minutes. Seventy-five (75)  $\mu$ g of Crotalus atrox venom then was added, and the incubation was continued for 3 minutes (15 minutes total). The reaction was stopped by addition of 200  $\mu$ L of activated charcoal (25 mg/mL suspension in 0.1 M NaH<sub>2</sub>PO<sub>4</sub>, pH 4). After centrifugation (750 X g for

- 97 -

3 minutes) to sediment the charcoal, a sample of the supernatant was taken for radioactivity determination in a scintillation counter and the PDE activity was calculated.

5

**Purification of PDE5 from *S. cerevisiae***

Cell pellets (29 g) were thawed on ice with an equal volume of Lysis Buffer (25 mM Tris HCl, pH 8, 5 mM MgCl<sub>2</sub>, 0.25 mM DTT, 1 mM benzamidine, and 10 μM ZnSO<sub>4</sub>). Cells were lysed in a Microfluidizer<sup>®</sup> (Microfluidics Corp.) using nitrogen at 20,000 psi. The lysate was centrifuged and filtered through 0.45 μm disposable filters. The filtrate was applied to a 150 mL column of Q SEPHAROSE<sup>®</sup> Fast-Flow (Pharmacia). The column was washed with 1.5 volumes of Buffer A (20 mM Bis-Tris Propane, pH 6.8, 1 mM MgCl<sub>2</sub>, 0.25 mM DTT, 10 μM ZnSO<sub>4</sub>) and eluted with a step gradient of 125 mM NaCl in Buffer A followed by a linear gradient of 125-1000 mM NaCl in Buffer A. Active fractions from the linear gradient were applied to a 180 mL hydroxyapatite column in Buffer B (20 mM Bis-Tris Propane (pH 6.8), 1 mM MgCl<sub>2</sub>, 0.25 mM DTT, 10 μM ZnSO<sub>4</sub>, and 250 mM KCl). After loading, the column was washed with 2 volumes of Buffer B and eluted with a linear gradient of 0-125 mM potassium phosphate in Buffer B. Active fractions were pooled, precipitated with 60% ammonium sulfate, and resuspended in Buffer C (20 mM Bis-Tris Propane, pH 6.8, 125 mM NaCl, 0.5 mM DTT, and 10 μM ZnSO<sub>4</sub>). The pool was applied to a 140 mL column of SEPHACRYL<sup>®</sup> S-300 HR and eluted with Buffer C.

- 98 -

Active fractions were diluted to 50% glycerol and stored at -20°C.

The resultant preparations were about 85% pure by SDS-PAGE. These preparations had specific activities of about 3  $\mu$ mol cGMP hydrolyzed per minute per milligram protein.

#### Inhibitory Effect on cGMP-PDE

cGMP-PDE activity of compounds of the present invention was measured using a one-step assay adapted from Wells et al., *Biochim. Biophys. Acta*, 384, 430 (1975). The reaction medium contained 50 mM Tris-HCl, pH 7.5, 5 mM magnesium acetate, 250  $\mu$ g/ml 5'-Nucleotidase, 1 mM EGTA, and 0.15  $\mu$ M 8-[H<sup>3</sup>]-cGMP. Unless otherwise indicated, the enzyme used was a human recombinant PDE5 (ICOS Corp., Bothell, Washington).

Compounds of the invention were dissolved in DMSO finally present at 2% in the assay. The incubation time was 30 minutes during which the total substrate conversion did not exceed 30%.

The IC<sub>50</sub> values for the compounds examined were determined from concentration-response curves typically using concentrations ranging from 10 nM to 10  $\mu$ M. Tests against other PDE enzymes using standard methodology showed that compounds of the invention are selective for the cGMP-specific PDE enzyme.

30

- 99 -

Biological Data

The compounds according to the present invention were typically found to exhibit an IC<sub>50</sub> value of less than 500 nM (i.e., 0.5  $\mu$ M). In vitro test data for representative compounds of the invention is given in the following table:

Table 1: <i>In vitro</i> Results (Examples)	
Example	PDE5 IC <sub>50</sub> ( $\mu$ M)
1	0.044
3	0.241
4 <sup>1)</sup>	0.2
5 <sup>1)</sup>	0.49
6	0.23
7 <sup>1)</sup>	1
8	.502
9	.261
11	.629
12	.03
13	.42

<sup>1)</sup> vs. bovine aorta

25

The intermediates also exhibited a low IC<sub>50</sub> value of less than 900 nM, typically less than 1  $\mu$ M (i.e., 1000 nM), and often less than 0.5  $\mu$ M, as illustrated in the following summary of *in vitro* test data.

30

- 100 -

Table 2: <i>In vitro</i> Results (Intermediates)	
Intermediate	PDE5 IC <sub>50</sub> ( $\mu$ M)
1a <sup>1)</sup>	0.85
1b <sup>1)</sup>	0.65
2 <sup>1)</sup>	0.2
3 <sup>1)</sup>	0.5
4 <sup>1)</sup>	0.8
5 <sup>1)</sup>	0.9
6	0.51
7a <sup>1)</sup>	0.2
7b <sup>1)</sup>	0.7
8	0.76
9a <sup>1)</sup>	0.47
9b	0.36
10	0.86
11	0.89
12	0.96
13a	0.27
13b	0.3
14	0.48
15a	0.29
15b	0.92
16a	0.53
16b	0.31
17a	0.38
17b	0.12
18	0.42
19	0.36

- 101 -

Table 2: <i>In vitro</i> Results (Intermediates)	
Intermediate	PDE5 IC <sub>50</sub> ( $\mu$ M)
20a	0.65
20b	0.34
21	0.68
22	0.68
5 23	0.8
24a	0.94
24b	0.92
25	0.59
26a	0.82
10 26b	0.54
27	0.83
28	0.22
29a	0.34
29b	0.23
15 30a	0.78
30b	0.42
31	0.43
32	0.33
33	0.86
20 34	0.23
35	0.16
36	0.09
37	0.62
38	0.77
25 39	0.65

<sup>1)</sup> vs. bovine aorta


- 102 -

Obviously, many modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and, therefore, only such limitations  
5 should be imposed as are indicated by the appended claims.

- 103 -

## WHAT IS CLAIMED IS:

1. A compound having a formula




wherein  $R^0$ , independently, is selected from the group consisting of halo,  $C_{1-6}$ alkyl, aryl, heteroaryl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{3-8}$ cycloalkylQ,  $C(=O)R^a$ ,  $OC(=O)R^a$ ,  $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^c$ ,  $C(=O)C_{1-4}$ alkyleneHet,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $NR^aR^b$ ,  $OC_{1-4}$ alkyleneHet,  $OC_{1-4}$ alkylene $OR^a$ ,  $OC_{1-4}$ alkylene $NR^aC(=O)OR^b$ ,  $NR^aR^b$ ,  $NR^aC_{1-4}$ alkylene $NR^aR^b$ ,  $NR^aC(=O)R^b$ ,  $NR^aC(=O)NR^aR^b$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $NR^a(SO_2C_{1-4}alkyl)$ , nitro, trifluoromethyl, trifluoromethoxy, cyano,  $SO_2NR^aR^b$ ,  $SO_2R^a$ ,  $SOR^a$ ,  $SR^a$ , and  $OSO_2CF_3$ ;



$R^1$  is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, an optionally substituted  $C_{3-8}$ cycloalkyl ring, an optionally substituted  $C_{3-8}$ heterocycloalkyl ring, an optionally substituted bicyclic ring




- 104 -




wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one to three heteroatoms selected from oxygen, sulfur, and nitrogen, hydrogen,  $C_{1-6}$ alkyl, aryl $C_{1-3}$ alkyl,  $C_{1-3}$ -alkylenearyl, halo $C_{1-6}$ alkyl,  $C_{1-4}$ alkyleneC(=O)OR<sup>a</sup>,  $C_{1-4}$ alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkenyl,  $C_{3-8}$ heterocycloalkenyl,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene-QR<sup>a</sup>,  $C_{2-6}$ alkenyleneQR<sup>a</sup>,  $C_{1-4}$ alkyleneQC $C_{1-4}$ alkyleneQR<sup>a</sup>,



- 105 -



and a spiro substituent having a structure



$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{2-6}$ alkenyl,  $C_{1-3}$ alkylenearyl, aryl $C_{1-3}$ alkyl,  $C(=O)R^a$ , aryl, heteroaryl,  $C(=O)R^a$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $SO_2R^a$ ,  $SO_2NR^aR^b$ ,  $S(=O)R^a$ ,  $S(=O)NR^aR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneOR<sup>a</sup>,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)C_{1-4}$ alkylenearyl,  $C(=O)C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkylenearyl substituted with one or more of  $SO_2NR^aR^b$ ,  $NR^aR^b$ ,  $C(=O)OR^a$ ,  $NR^aSO_2CF_3$ , CN,  $NO_2$ ,  $C(=O)R^a$ ,  $OR^a$ ,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>, and  $OC_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneC(=O)Het,  $C_{1-4}$ alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>C(=O) $R^a$ ,  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneC(=O)OR<sup>a</sup>, and  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>;


- 106 -

$R^3$  is selected from the group consisting of  $C(=O)R^b$ ,  $C(=O)OR^b$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $C(=O)Het$ ,  $C(=O)NR^aC_{1-4}alkyleneOR^a$ ,  $C(=O)NR^aC_{1-4}alkyleneHet$ ,  $C(=O)C_{1-4}alkylenearyl$ ,  $C(=O)C_{1-4}alkyleneheteroaryl$ ,  $C(=O)NR^aC_{1-4}alkylenearyl$ ,  $C(=O)NR^aC_{1-4}alkyleneC_{3-8}cycloalkyl$ ,  $C(=O)NR^bSO_2R^c$ ,  $C(=O)NR^aC_{1-4}alkyleneOC_{1-6}alkyl$ ,  $C(=O)NR^aC_{1-4}alkyleneheteroaryl$ ,  $NR^aR^c$ ,  $NR^aC(=O)R^b$ ,  $NR^aC(=O)NR^aR^c$ ,  $NR^a(SO_2C_{1-4}alkyl)$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $OR^a$ ,  $NR^aC(=O)C_{1-4}alkyleneN(R^b)_2$ ,  $NR^aC(=O)C_{1-4}alkyleneC(=O)OR^a$ ,  $NR^aC(=O)C_{1-3}alkylenearyl$ ,  $NR^aC(=O)C_{1-3}alkyleneC_{3-8}heterocycloalkyl$ ,  $NR^aC(=O)C_{1-3}alkyleneHet$ , and  $C(=O)NR^aSO_2R^b$ ;

$R^4$  is selected from the group consisting of hydrogen,  $C_{1-6}alkyl$ , aryl, heteroaryl, aryl $C_{1-3}alkyl$ ,  $C_{1-3}alkylenearyl$ ,  $C_{1-3}alkyleneHet$ ,  $C_{3-8}cycloalkyl$ , and  $C_{3-8}heterocycloalkyl$ ;

$X$  is selected from the group consisting of  $C(=O)$ ,  $(CH_2)_tC(=O)$ ,  $C(=O)C\equiv C$ ,  $C(=O)C(R^a)=C(R^a)$ ,  $C(=S)$ ,  $SO$ ,  $SO_2$ ,  $SO_2C(R^a)=CR^a$ ,  $CR^aR^b$ ,  $CR^a=CR^a$ ,  $C(=O)NR^a$ , and  $C(=N-OR^a)$ ;

$Y$  is selected from the group consisting of  $R^a$ ,  $R^d$ ,  $(CH_2)_nC(=O)R^c$ ,  $N(R^b)(CH_2)_nR^c$ ,  $O(CH_2)_nR^c$ ,  $N(R^b)C(=O)R^c$ ,  $C(=O)N(R^a)(R^c)$ ,  $N(R^a)C(=O)R^c$ ,



and  $N(R^a)SO_2R^c$ ;

- 107 -

R<sup>a</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-6</sub>cycloalkyl, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, heteroaryl, heteroarylC<sub>1-3</sub>alkyl, and C<sub>1-3</sub>alkyleneheteroaryl;

R<sup>b</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and heteroaryl;

R<sup>c</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>1-6</sub>alkylenearyl, C<sub>1-6</sub>alkyleneHet, haloC<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>heterocycloalkyl, Het, C<sub>1-3</sub>alkyleneheteroaryl, C<sub>1-6</sub>alkyleneC(=O)OR<sup>a</sup>, and C<sub>1-3</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl;

or R<sup>a</sup> and R<sup>c</sup> are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;

R<sup>d</sup> is a 5- or 6-membered ring or a bicyclic fused ring system, saturated or partially or fully unsaturated, comprising carbon atoms and optionally one to three heteroatoms selected from oxygen, sulfur, and nitrogen, and optionally substituted with one or more R<sup>e</sup> or R<sup>f</sup>;

R<sup>e</sup> is selected from the group consisting of  
nitro,  
trifluoromethyl,  
trifluoromethoxy,  
halogen,  
cyano,  
Het,  
C<sub>1-6</sub>alkyl,  
C<sub>1-6</sub>alkyleneOR<sup>a</sup>,  
C(=O)R<sup>a</sup>,

- 108 -

$\text{OC}(=\text{O})\text{R}^a$ ,  
 $\text{C}(=\text{O})\text{OR}^a$ ,  
 $\text{C}_{1-4}\text{alkyleneHet}$ ,  
 $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ,  
 $\text{OC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ,  
 $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ,  
 $\text{C}(=\text{O})\text{NR}^a\text{SO}_2\text{R}^f$ ,  
 $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneHet}$ ,  
 $\text{C}_{1-4}\text{alkyleneNR}^a\text{R}^b$ ,  
 $\text{C}_{2-6}\text{alkenyleneNR}^a\text{R}^b$ ,  
 $\text{C}(=\text{O})\text{NR}^a\text{R}^b$ ,  
 $\text{C}(=\text{O})\text{NR}^a\text{R}^g$ ,  
 $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneOR}^b$ ,  
 $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneHet}$ ,  
 $\text{OR}^a$ ,  
 $\text{OC}_{2-4}\text{alkyleneNR}^a\text{R}^b$ ,  
 $\text{OC}_{1-4}\text{alkyleneCH}(\text{OR}^a)\text{CH}_2\text{-NR}^a\text{R}^b$ ,  
 $\text{OC}_{1-4}\text{alkyleneHet}$ ,  
 $\text{OC}_{2-4}\text{alkyleneOR}^a$ ,  
 $\text{OC}_{2-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{OR}^b$ ,  
 $\text{NR}^a\text{R}^b$ ,  
 $\text{NR}^a\text{C}_{1-4}\text{alkyleneNR}^a\text{R}^b$ ,  
 $\text{NR}^a\text{C}(=\text{O})\text{R}^b$ ,  
 $\text{NR}^a\text{C}(=\text{O})\text{NR}^a\text{R}^b$ ,  
 $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,  
 $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ ,  
 $\text{SO}_2\text{NR}^a\text{R}^b$ ,  
 and  $\text{OSO}_2\text{trifluoromethyl}$ ;

$\text{R}^f$  is selected from the group consisting of  
 hydrogen, halogen,  $\text{OR}^a$ ,  $\text{C}_{1-6}\text{alkyl}$ , nitro, and  $\text{NR}^a\text{R}^b$ ;

or  $\text{R}^e$  and  $\text{R}^f$  are taken together to form a  
 3- or 4-membered alkylene or alkenylene chain

- 109 -

component of a 5- or 6-membered ring, optionally containing at least one heteroatom;

R<sup>g</sup> is phenyl or C<sub>4-6</sub>cycloalkyl, optionally substituted with one or more halogen, C(=O)OR<sup>a</sup>, or OR<sup>a</sup>;

Q is O, S, or NR<sup>h</sup>;

B is O, S, or NR<sup>h</sup>;

C is O, S, or NR<sup>a</sup>;

D is CR<sup>a</sup> or N;

E is CR<sup>a</sup>, C(R<sup>a</sup>)<sub>2</sub>, or NR<sup>h</sup>; and

R<sup>h</sup> is null or is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and C<sub>1-3</sub>alkyleneheteroaryl;

Het represents a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>a</sup>;

n is 0 or 1;


q is 0, 1, 2, 3, or 4;

t is 1, 2, 3, or 4;

and pharmaceutically acceptable salts and solvates thereof.

- 110 -

2. The compound of claim 1 represented by the formula






and pharmaceutically acceptable salts and hydrates thereof.


3. The compound of claim 1 wherein q is 0, or R<sup>0</sup> is selected from the group consisting of aryl, Het, OR<sup>a</sup>, C(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkylQ, C(=O)NR<sup>a</sup>R<sup>b</sup>, and C(=O)NR<sup>a</sup>R<sup>c</sup>.

- 111 -

4. The compound of claim 1 wherein  $R^1$  is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl,  $C_{1-4}$ alkyleneQR<sup>a</sup>,  $C_{1-4}$ alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkenyl,  $C_{1-6}$ alkyl,




, and






- 112 -

5. The compound of claim 1 wherein  $R^1$  is the optionally substituted bicyclic ring system




6. The compound of claim 5 wherein  $R^1$  is




and wherein  $p$  is an integer 1 or 2, and  $G$ , independently, are  $C(R^a)_2$ , O, S, or  $NR^a$ .

- 113 -


7. The compound of claim 1 wherein R<sup>1</sup> is selected from the group consisting of




,



,







,








,


- 114 -




- 115 -



- 116 -

-CH<sub>2</sub>OR<sup>a</sup>, -CH<sub>2</sub>OCH<sub>2</sub>OR<sup>a</sup>,


, and




8. The compound of claim 1 wherein R<sup>2</sup> is selected from the group consisting of hydrogen, aryl, heteroaryl, OR<sup>a</sup>, NR<sup>a</sup>R<sup>b</sup>, NR<sup>a</sup>R<sup>c</sup>, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkylenearyl, C<sub>1-4</sub>alkyleneC(=O)C<sub>1-4</sub>alkylenearyl, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>, C<sub>1-4</sub>alkyleneC(=O)NR<sup>a</sup>R<sup>c</sup>, C<sub>1-4</sub>alkyleneC(=O)Het, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>c</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>C(=O)R<sup>a</sup>, and C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneOR<sup>a</sup>.

- 117 -

9. The compound of claim 8 wherein  $R^2$  is selected from the group consisting of hydrogen;  $C_{1-4}$ -alkyleneheteroaryl, wherein the heteroaryl group is selected from the group consisting of benzimidazole, a triazole, and imidazole;  $C_{1-4}$ alkyleneHet, wherein Het is selected from the group consisting of piperazine, morpholine, pyrrolidine, pyrrolidone, tetrahydrofuran, piperidine,



, and




;

$C_{1-4}$ alkylene $C_6H_5$ , optionally substituted with one to three groups selected from the group consisting of  $C(=O)OR^a$ ,  $NR^aR^b$ ,  $NR^aSO_2CF_3$ ,  $SO_2NR^aR^b$ , CN,  $OR^a$ ,  $C(=O)R^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ , nitro,  $OC_{1-4}$ alkylenearyl, and  $OC_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)$ benzyl;  $C_{1-4}$ alkylene $C(=O)OR^a$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^c$ ;  $C_{1-4}$ alkyleneHet;  $NR^aR^b$ ; OH;  $OC_{1-4}$ alkyl;  $C_6H_5$ ;  $C_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $OR^a$ ;  $C_{1-4}$ alkylene $NHC(=O)R^a$ ; and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ .

- 118 -

10. The compound of claim 8 wherein  $R^2$  is selected from the group consisting of  $C_{1-6}$ alkyl,  $C(=O)OR^a$ ,  $C(=O)R^a$ , hydrogen,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)NR^aR^c$ , aryl, and heteroaryl.

11. The compound of claim 1 wherein  $R^3$  is selected from the group consisting of  $C(=O)OR^a$ ,  $C(=O)R^a$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OC_{1-6}$ alkyl,  $C(=O)NR^aC_{1-4}$ alkylene $C_{3-8}$ cycloalkyl,  $C(=O)Het$ ,



$C(=O)NR^aC_{1-4}$ alkyleneheteroaryl,  $C(=O)NR^aC_{1-4}$ alkylene-aryl,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)NR^aR^c$ , and  $C(=S)NR^aR^c$ .





12. The compound of claim 1 wherein  $R^4$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, and heteroaryl.

13. The compound of claim 1 wherein  $n$  is 0, or  $X$  is selected from the group consisting of  $C(=O)$ ,  $C(=O)CR^a=CR^a$ ,  $(CH_2)_tC=O$ ,  $C(=S)$  and  $C(=N-OR^a)$ .

14. The compound of claim 1 wherein  $Y$  is selected from the group consisting of  $R^a$ ,  $R^d$ ,  $N(R^b)(CH_2)_nR^c$ ,  $N(R^b)C(=O)R^c$ , and  $N(R^a)C(=O)R^c$ .




- 119 -

15. The compound of claim 1 wherein q is 0, or R<sup>0</sup> is selected from the group consisting of halo, methyl, trifluoromethyl, and trifluoromethyl; R<sup>1</sup> is selected from the group consisting of







- 120 -



- 121 -




, and




;

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C(=O)NR^aR^c$ , and  $C_{1-4}$ alkyleneHet;  $R^3$  is selected from the group consisting of  $C(=O)OC_2H_5$ ,  $C(=O)OCH_3$ ,  $C(=O)NHCH_2C_6H_5$ ,  $C(=O)NH(CH_2)_2C_6H_5$ ,  $C(=O)NHC_6H_5$ ,  $C(=O)NH_2$ ,  $C(=O)N(CH_3)_2$ ,  $C(=S)N(CH_3)_2$ ,  $C(=O)NH(CH_2)_2CH_3$ ,  $C(=O)N(CH_2)_3CH_3$ ,  $C(=O)NHCH_3$ ,  $C(=O)NHCH(CH_3)_2$ ,  $C(=O)NH(CH_2)_3OCH_3$ ,








,




,


- 122 -



- 123 -




,




,



,








,





,


- 124 -



- 125 -



, and






;

and  $R^4$  is hydrogen or  $C_{1-6}$ alkyl.




16. The compound of claim 1 wherein  $q$  is 0,  $R^2$  is hydrogen, and  $R^4$  is hydrogen or methyl.

- 126 -

17. A compound selected from the group consisting of






- 127 -









- 128 -



- 129 -



- 130 -



and pharmaceutically acceptable salts and solvates thereof.

18. A pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

19. A method of treating a male or female animal in the treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit comprising treating said animal with an effective amount of a pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

20. The method of claim 19 wherein the condition is male erectile dysfunction.

21. The method of claim 20 wherein the treatment is an oral treatment.

- 131 -

22. The method of claim 19 wherein the condition is female arousal disorder.

23. The method of claim 22 wherein the treatment is an oral treatment.

24. The method of claim 19 wherein the condition is selected from the group consisting of stable angina, unstable angina, variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, malignant hypertension, pheochromocytoma, acute respiratory distress syndrome, congestive heart failure, acute renal failure, chronic renal failure, atherosclerosis, a condition of reduced blood vessel patency, a peripheral vascular disease, a vascular disorder, thrombocythemia, an inflammatory disease, myocardial infarction, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, peptic ulcer, a gut motility disorder, postpercutaneous transluminal coronary angioplasty, carotid angioplasty, post-bypass surgery graft stenosis, osteoporosis, preterm labor, benign prostatic hypertrophy, and irritable bowel syndrome.

25. A method of treating a condition where inhibition of a cGMP-specific PDE is of therapeutic benefit, in a human or a nonhuman animal body, comprising administering to said body a therapeutically effective amount of a compound of claim 1.


- 132 -

26. A method for the curative or prophylactic treatment of male erectile dysfunction or female arousal disorder, comprising administration of an effective dose of a compound of claim 1, and pharmaceutically acceptable salts and solvates thereof, to an animal.

27. Use of a compound of claim 1 for the manufacture of a medicament for the curative or prophylactic treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit.

- 133 -


28. A compound having a formula






wherein  $R^0$ , independently, is selected from the group consisting of halo,  $C_{1-6}$ alkyl, aryl, heteroaryl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{3-8}$ cycloalkylQ,  $C(=O)R^a$ ,  $OC(=O)R^a$ ,  $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^c$ ,  $C(=O)C_{1-4}$ alkyleneHet,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $NR^aR^b$ ,  $OC_{1-4}$ alkyleneHet,  $OC_{1-4}$ alkylene $OR^a$ ,  $OC_{1-4}$ alkylene $NR^aC(=O)OR^b$ ,  $NR^aR^b$ ,  $NR^aC_{1-4}$ alkylene $NR^aR^b$ ,  $NR^aC(=O)R^b$ ,  $NR^aC(=O)NR^aR^b$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $NR^a(SO_2C_{1-4}alkyl)$ , nitro, trifluoromethyl, trifluoromethoxy, cyano,  $SO_2NR^aR^b$ ,  $SO_2R^a$ ,  $SOR^a$ ,  $SR^a$ , and  $OSO_2CF_3$ ;

$R^1$  is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl, an optionally substituted  $C_{3-8}$ cycloalkyl ring, an optionally substituted  $C_{3-8}$ heterocycloalkyl ring, an optionally substituted bicyclic ring

- 134 -




wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one to three heteroatoms selected from oxygen, sulfur, and nitrogen, hydrogen, C<sub>1-6</sub>alkyl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>-alkylenearyl, haloC<sub>1-6</sub>alkyl, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkenyl, C<sub>3-8</sub>heterocycloalkenyl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkylene-QR<sup>a</sup>, C<sub>2-6</sub>alkenyleneQR<sup>a</sup>, C<sub>1-4</sub>alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>,



- 135 -

and a spiro substituent having a structure



;

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{2-6}$ alkenyl,  $C_{1-3}$ alkylenearyl, aryl $C_{1-3}$ alkyl,  $C(=O)R^a$ , aryl, heteroaryl,  $C(=O)R^a$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $SO_2R^a$ ,  $SO_2NR^aR^b$ ,  $S(=O)R^a$ ,  $S(=O)NR^aR^b$ ,  $C(=O)NR^aC_{1-4}$ alkyleneOR<sup>a</sup>,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)C_{1-4}$ alkylenearyl,  $C(=O)C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkylenearyl substituted with one or more of  $SO_2NR^aR^b$ ,  $NR^aR^b$ ,  $C(=O)OR^a$ ,  $NR^aSO_2CF_3$ , CN,  $NO_2$ ,  $C(=O)R^a$ ,  $OR^a$ ,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>, and  $OC_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneC(=O)Het,  $C_{1-4}$ alkyleneC(=O)NR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>C(=O) $R^a$ ,  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneOR<sup>a</sup>,  $C_{1-4}$ alkyleneNR<sup>a</sup>R<sup>b</sup>,  $C_{1-4}$ alkyleneC(=O)OR<sup>a</sup>, and  $C_{1-4}$ alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>;

$R^3$  is selected from the group consisting of  $C(=O)R^b$ ,  $C(=O)OR^b$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $C(=S)NR^aR^b$ ,  $C(=S)NR^aR^c$ ,  $C(=O)Het$ ,  $C(=O)NR^aC_{1-4}$ alkyleneOR<sup>a</sup>,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)C_{1-4}$ alkylenearyl,  $C(=O)C_{1-4}$ alkyleneheteroaryl,  $C(=O)NR^aC_{1-4}$ alkylenearyl,  $C(=O)NR^aC_{1-4}$ alkyleneC<sub>3-8</sub>cycloalkyl,  $C(=O)NR^bSO_2R^c$ ,  $C(=O)-$



- 136 -

NR<sup>a</sup>C<sub>1-4</sub>alkyleneOC<sub>1-6</sub>alkyl, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneheteroaryl, NR<sup>a</sup>R<sup>c</sup>, NR<sup>a</sup>C(=O)R<sup>b</sup>, NR<sup>a</sup>C(=O)NR<sup>a</sup>R<sup>c</sup>, NR<sup>a</sup>(SO<sub>2</sub>C<sub>1-4</sub>alkyl), N(SO<sub>2</sub>C<sub>1-4</sub>alkyl)<sub>2</sub>; OR<sup>a</sup>, NR<sup>a</sup>C(=O)C<sub>1-4</sub>alkyleneN(R<sup>b</sup>)<sub>2</sub>, NR<sup>a</sup>C(=O)C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, NR<sup>a</sup>(C=O)C<sub>1-3</sub>alkylenearyl, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkyleneHet, and C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>b</sup>;

R<sup>a</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, C<sub>1-3</sub>alkyleneHet, C<sub>3-8</sub>cycloalkyl, and C<sub>3-8</sub>heterocycloalkyl;

R<sup>a</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-6</sub>cycloalkyl, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, heteroaryl, heteroarylC<sub>1-3</sub>alkyl, and C<sub>1-3</sub>alkyleneheteroaryl;

R<sup>b</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and heteroaryl;

R<sup>c</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>1-6</sub>alkylenearyl, C<sub>1-6</sub>alkyleneHet, haloC<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>heterocycloalkyl, Het, C<sub>1-3</sub>alkyleneheteroaryl, C<sub>1-6</sub>alkyleneC(=O)OR<sup>a</sup>, and C<sub>1-3</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl;

or R<sup>a</sup> and R<sup>c</sup> are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;

Q is O, S, or NR<sup>h</sup>;

B is O, S, or NR<sup>h</sup>;

C is O, S, or NR<sup>a</sup>;

D is CR<sup>a</sup> or N;

E is CR<sup>a</sup>, C(R<sup>a</sup>)<sub>2</sub>, or NR<sup>h</sup>; and

- 137 -

R<sup>n</sup> is null or is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and C<sub>1-3</sub>alkyleneheteroaryl;

Het represents a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>a</sup>;

n is 0 or 1;


q is 0, 1, 2, 3, or 4;

t is 1, 2, 3, or 4;

and pharmaceutically acceptable salts and solvates thereof.

- 138 -

29. The compound of claim 28 represented by the formula






and pharmaceutically acceptable salts and hydrates thereof.


30. The compound of claim 28 wherein q is 0, or R<sup>0</sup> is selected from the group consisting of aryl, Het, OR<sup>a</sup>, C(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkylQ, C(=O)NR<sup>a</sup>R<sup>b</sup>, and C(=O)NR<sup>a</sup>R<sup>c</sup>.

- 139 -

31. The compound of claim 28 wherein  $R^1$  is selected from the group consisting of optionally substituted aryl, optionally substituted heteroaryl,  $C_{1-4}$ alkyleneQR<sup>a</sup>,  $C_{1-4}$ alkyleneQC<sub>1-4</sub>alkyleneQR<sup>a</sup>,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkenyl,  $C_{1-6}$ alkyl,




, and




- 140 -

32. The compound of claim 28 wherein  $R^1$  is the optionally substituted bicyclic ring system








33. The compound of claim 32 wherein  $R^1$  is




and wherein  $p$  is an integer 1 or 2, and  $G$ , independently, are  $C(R^a)_2$ ,  $O$ ,  $S$ , or  $NR^a$ .

- 141 -


34. The compound of claim 28 wherein R<sup>1</sup> is selected from the group consisting of




- 142 -




,




,



,




,




,

- 143 -




,




,

 $-\text{CH}_2\text{OR}^a$ ,  $-\text{CH}_2\text{OCH}_2\text{OR}^a$ 

,



, and






- 144 -


35. The compound of claim 28 wherein  $R^2$  is selected from the group consisting of aryl, hydrogen, heteroaryl,  $OR^a$ ,  $NR^aR^b$ ,  $NR^aR^c$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $OR^a$ ,  $C_{1-4}$ alkyleneC(=O) $NR^aR^b$ ,  $C_{1-4}$ alkyleneC(=O) $NR^aR^c$ ,  $C_{1-4}$ alkyleneC(=O)Het,  $C_{1-4}$ alkylene $NR^aR^b$ ,  $C_{1-4}$ alkylene $NR^aR^c$ ,  $C_{1-4}$ alkylene $NR^aC(=O)R^a$ , and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ .

- 145 -

36. The compound of claim 28 wherein  $R^2$  is selected from the group consisting of hydrogen;  $C_{1-4}$ -alkyleneheteroaryl, wherein the heteroaryl group is selected from the group consisting of benzimidazole, a triazole, and imidazole;  $C_{1-4}$ alkyleneHet, wherein Het is selected from the group consisting of piperazine, morpholine, pyrrolidine, pyrrolidone, tetrahydrofuran, piperidine,



, and




;

$C_{1-4}$ alkylene $C_6H_5$ , optionally substituted with one to three groups selected from the group consisting of  $C(=O)OR^a$ ,  $NR^aR^b$ ,  $NR^aSO_2CF_3$ ,  $SO_2NR^aR^b$ , CN,  $OR^a$ ,  $C(=O)R^a$ ,  $C_{1-4}$ alkylene $NR^aR^b$ , nitro,  $OC_{1-4}$ alkylenearyl, and  $OC_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)$ benzyl;  $C_{1-4}$ alkylene $C(=O)OR^a$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^b$ ;  $C_{1-4}$ alkylene $C(=O)NR^aR^c$ ;  $C_{1-4}$ alkyleneHet;  $NR^aR^b$ ; OH;  $OC_{1-4}$ alkyl;  $C_6H_5$ ;  $C_{1-4}$ alkylene $NR^aR^b$ ;  $C_{1-4}$ alkylene $OR^a$ ;  $C_{1-4}$ alkyleneNHC(=O) $R^a$ ; and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ .

- 146 -

37. The compound of claim 28 wherein  $R^2$  is selected from the group consisting of  $C_{1-6}$ alkyl,  $C(=O)OR^a$ ,  $C(=O)R^a$ , hydrogen,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)NR^aR^c$ , aryl, and heteroaryl.

38. The compound of claim 28 wherein  $R^3$  is selected from the group consisting of  $C(=O)OR^a$ ,  $C(=O)R^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneOC $_{1-6}$ alkyl,  $C(=O)NR^aC_{1-4}$ alkyleneC $_{3-9}$ cycloalkyl,  $C(=O)Het$ ,







$C(=O)NR^aC_{1-4}$ alkyleneheteroaryl,  $C(=O)NR^aC_{1-4}$ alkylene-aryl,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)NR^aR^c$ , and  $C(=S)NR^aR^c$ .





39. The compound of claim 28 wherein  $R^4$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, and heteroaryl.

- 147 -

40. The compound of claim 1 wherein q is 0, or R<sup>0</sup> is selected from the group consisting of halo, methyl, trifluoromethyl, and trifluoromethyl; R<sup>1</sup> is selected from the group consisting of




- 148 -




, and

- 149 -




;

$R^2$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C(=O)NR^aR^c$ , and  $C_{1-4}$ allyleneHet;  $R^3$  is selected from the group consisting of  $C(=O)OC_2H_5$ ,  $C(=O)OCH_3$ ,  $C(=O)NHCH_2C_6H_5$ ,  $C(=O)NH(CH_2)_2C_6H_5$ ,  $C(=O)NHC_6H_5$ ,  $C(=O)NH_2$ ,  $C(=O)N(CH_3)_2$ ,  $C(=S)N(CH_3)_2$ ,  $C(=O)NH(CH_2)_2CH_3$ ,  $C(=O)N(CH_2)_3CH_3$ ,  $C(=O)NHCH_3$ ,  $C(=O)NHCH(CH_3)_2$ ,  $C(=O)NH(CH_2)_3OCH_3$ ,








,




,


- 150 -




- 151 -




,




,




,



,








,




,




- 152 -



- 153 -



, and






;

and  $\text{R}^4$  is hydrogen and  $\text{C}_{1-6}$ alkyl.




41. The compound of claim 28 wherein  $q$  is 0,  $\text{R}^2$  is hydrogen, and  $\text{R}^4$  is hydrogen or methyl.

- 154 -




42. A compound selected from the group consisting of






- 155 -






- 156 -






- 157 -



- 158 -






- 159 -











- 160 -






- 161 -






- 162 -






- 163 -






- 164 -






- 165 -



- 166 -






- 167 -











- 168 -



- 169 -



- 170 -



and pharmaceutically acceptable salts and solvates thereof.

- 171 -

43. A pharmaceutical composition comprising a compound of claim 28, together with a pharmaceutically acceptable diluent or carrier.

44. A method of treating a male or female animal in the treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit comprising treating said animal with an effective amount of a pharmaceutical composition comprising a compound of claim 28, together with a pharmaceutically acceptable diluent or carrier.

45. The method of claim 44 wherein the condition is male erectile dysfunction.

46. The method of claim 44 wherein the condition is female arousal disorder.

47. The method of claim 45 wherein the treatment is an oral treatment.

48. The method of claim 46 wherein the treatment is an oral treatment.

- 172 -

49. The method of claim 44 wherein the condition is selected from the group consisting of stable angina, unstable angina, variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, malignant hypertension, pheochromocytoma, acute respiratory distress syndrome, congestive heart failure, acute renal failure, chronic renal failure, atherosclerosis, a condition of reduced blood vessel patency, a peripheral vascular disease, a vascular disorder, thrombocythemia, an inflammatory disease, myocardial infarction, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, peptic ulcer, a gut motility disorder, postpercutaneous transluminal coronary angioplasty, carotid angioplasty, post-bypass surgery graft stenosis, osteoporosis, preterm labor, benign prostatic hypertrophy, and irritable bowel syndrome.

50. A method of treating a condition where inhibition of a cGMP-specific PDE is of therapeutic benefit, in a human or a nonhuman animal body, comprising administering to said body a therapeutically effective amount of a compound of claim 28.

51. A method for the curative or prophylactic treatment of male erectile dysfunction or female arousal disorder, comprising administration of an effective dose of a compound of claim 28, and pharmaceutically acceptable salts and solvates thereof, to an animal.

- 173 -

52. Use of a compound of claim 28 for the manufacture of a medicament for the curative or prophylactic treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 02/10367

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D471/04 A61K31/435 A61P15/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 01 87038 A (ORTHO MCNEIL PHARM INC) 22 November 2001 (2001-11-22) claim 1; tables 1-5 ---	1-4, 28-31
P, X	WO 02 28865 A (DAUGAN ALAIN CLAUDE MARIE ;GOSIMINI ROMAIN LUC MARIE (FR); ORME MA) 11 April 2002 (2002-04-11) Intermediate compounds 1-4 claim 1 ---	1-52
P, X	WO 02 36593 A (ORME MARK W ;LILLY ICOS LLC (US); SCHULTZE LISA M (US); SAWYER JAS) 10 May 2002 (2002-05-10) claim 1 --- -/--	1-52

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

24 July 2002

Date of mailing of the international search report

12/08/2002

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Baston, E

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/10367

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 94347 A (DAUGAN ALAIN CLAUDE MARIE ;BOMBRUN AGNES (FR); ORME MARK W (US); L) 13 December 2001 (2001-12-13) claim 1 ---	1-52
X	EP 0 466 548 A (ADIR) 15 January 1992 (1992-01-15) claim 1 ---	1-4, 28-31
X	US 6 069 150 A (MENTA ERNESTO ET AL) 30 May 2000 (2000-05-30) claim 1 ---	1-52
X	US 4 336 260 A (PAYNE ALAN J ET AL) 22 June 1982 (1982-06-22) claim 1 ---	1-52
Y	WO 96 32003 A (DAUGAN ALAIN CLAUDE MARIE ;GLAXO WELLCOME LAB SA (FR); LABAUDINIER) 17 October 1996 (1996-10-17) claim 1 ---	1-52
Y	WO 01 08686 A (KRAL MARTHA A ;ANDERSON NEIL R (US); LILLY ICOS LLC (US); OREN PET) 8 February 2001 (2001-02-08) claim 1 ---	1-52
Y	WO 00 15639 A (GELLIBERT FRANCOISE ;BOMBRUN AGNES (FR); ICOS CORP (US)) 23 March 2000 (2000-03-23) claim 1; examples 1-27 ---	1-52
Y	US 6 043 252 A (BOMBRUN AGNES) 28 March 2000 (2000-03-28) claim 1 ---	1-52
Y	WO 97 43287 A (BOMBRUN AGNES ;ICOS CORP (US)) 20 November 1997 (1997-11-20) claim 1 ---	1-52
X	KUMAR, S. ET AL.: "Possible Anthelmintic Agents" INDIAN JOURNAL OF CHEMISTRY, vol. 22B, 1983, pages 54-59, XP002190316 table 1 ---	1-52
X	SRIVASTAVA, S.K. ET AL.: "Potent 1,3-Disubstituted-9-Hpyrido'3,4-b!indoles as New Lead Compounds in Antifilarial Chemotherapy" BIOORGANIC & MEDICINAL CHEMISTRY, vol. 7, 1999, pages 1223-1236, XP002207274 table 4 ---	1-52
	---	
	-/--	



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/10367

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ISHIDA A ET AL: "A NEW METHOD FOR THE PREPARATION OF 3,4-DIHYDRO- AND 1,2,3,4-TETRAHYDRO-BETA-CARBOLINES" CHEMICAL AND PHARMACEUTICAL BULLETIN, PHARMACEUTICAL SOCIETY OF JAPAN. TOKYO, JP, vol. 33, no. 8, 1985, pages 3237-3249, XP001019167 ISSN: 0009-2363 examples 5,6 -----	1-52

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/10367

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 0187038	A	22-11-2001	AU 6116701 A		26-11-2001
			WO 0187038 A2		22-11-2001
			US 2002010189 A1		24-01-2002
WO 0228865	A	11-04-2002	WO 0228865 A2		11-04-2002
WO 0236593	A	10-05-2002	WO 0236593 A1		10-05-2002
WO 0194347	A	13-12-2001	AU 6806501 A		17-12-2001
			WO 0194347 A1		13-12-2001
EP 0466548	A	15-01-1992	FR 2663935 A1		03-01-1992
			AU 636232 B2		22-04-1993
			AU 7929391 A		02-01-1992
			CA 2045450 A1		28-12-1991
			EP 0466548 A1		15-01-1992
			IE 912225 A1		01-01-1992
			JP 4243879 A		31-08-1992
			NZ 238709 A		26-05-1992
			OA 9374 A		15-09-1992
			PT 98101 A		31-03-1992
			ZA 9104970 A		29-04-1992
US 6069150	A	30-05-2000	IT MI960664 A1		06-10-1997
			IT MI962241 A1		29-04-1998
			AT 212549 T		15-02-2002
			AU 710079 B2		16-09-1999
			AU 2506997 A		29-10-1997
			BR 9708480 A		13-04-1999
			DE 69710182 D1		14-03-2002
			DK 891187 T3		29-04-2002
			EP 0891187 A1		20-01-1999
			JP 2000508302 T		04-07-2000
			CA 2250898 A1		16-10-1997
			CN 1215333 A		28-04-1999
			WO 9737658 A1		16-10-1997
			TR 9801970 T2		23-10-2000
US 4336260	A	22-06-1982	NONE		
WO 9632003	A	17-10-1996	AT 211139 T		15-01-2002
			AU 702548 B2		25-02-1999
			AU 6613896 A		30-10-1996
			BR 9609503 A		08-03-2000
			CA 2226759 A1		17-10-1996
			CN 1195350 A		07-10-1998
			DE 69618231 D1		31-01-2002
			WO 9632003 A2		17-10-1996
			EP 0859778 A2		26-08-1998
			JP 11509517 T		24-08-1999
			US 6143757 A		07-11-2000
			US 6218400 B1		17-04-2001
			US 6001847 A		14-12-1999
WO 0108686	A	08-02-2001	AU 4490900 A		19-02-2001
			AU 4491200 A		19-02-2001
			BR 0012863 A		16-04-2002
			EP 1200090 A1		02-05-2002

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 02/10367

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0108686	A	EP 1200091 A1 NO 20020532 A WO 0108686 A1 WO 0108687 A1	02-05-2002 26-03-2002 08-02-2001 08-02-2001
WO 0015639	A	23-03-2000 WO 0015639 A1 AU 1025899 A BR 9816018 A EP 1114048 A1	23-03-2000 03-04-2000 05-06-2001 11-07-2001
US 6043252	A	28-03-2000 AT 215950 T AU 711885 B2 AU 2891097 A BG 102879 A BR 9709230 A DE 69711882 D1 EE 9800390 A EP 0912567 A1 JP 2000513717 T MD 980248 A NO 985222 A SK 154498 A3 US 6117881 A	15-04-2002 21-10-1999 05-12-1997 30-09-1999 10-08-1999 16-05-2002 15-06-1999 06-05-1999 17-10-2000 31-10-2000 11-01-1999 18-01-2000 12-09-2000
WO 9743287	A	20-11-1997 AT 215950 T AU 711885 B2 AU 2891097 A BG 102879 A BR 9709230 A CN 1218471 A ,B CZ 9803585 A3 DE 69711882 D1 EE 9800390 A WO 9743287 A1 EP 0912567 A1 HU 9901478 A2 JP 2000513717 T MD 980248 A NO 985222 A PL 329875 A1 SK 154498 A3 TR 9802282 T2 US 6117881 A US 6306870 B1	15-04-2002 21-10-1999 05-12-1997 30-09-1999 10-08-1999 02-06-1999 17-03-1999 16-05-2002 15-06-1999 20-11-1997 06-05-1999 30-08-1999 17-10-2000 31-10-2000 11-01-1999 12-04-1999 18-01-2000 22-03-1999 12-09-2000 23-10-2001